

No books, no notes. Calculators are allowed.

Show all your work in your bluebook.

Show all your work in your bluebook. Start each problem on a new page.

1. (30 Points) Let

$$f(x) = -\frac{3}{2}x^2 + \frac{5}{2}x + 1.$$

Notice that $f(0) = 1$, $f(1) = 2$, and $f(2) = 0$ is a period-3 orbit.


- a. Is the orbit $\mathcal{O}_f^+(0)$ attracting or repelling?
- b. What is the Lyapunov exponent of $x_0 = 0$?

2. (50 Points) Let

$$f(x) = x^3 - \frac{9}{16}x.$$

Notice that

$$f'(x) = 3x^2 - \frac{9}{16} \geq -\frac{9}{16} > -1.$$



- a. Find the fixed points and determine their stability type as attracting or repelling.
- b. Find the critical points, where $f'(x) = 0$. Call the critical points x_c^\pm .
- c. Show that for x in $[x_c^-, x_c^+]$, $|f'(x)| < 1$, so by the Mean Value Theorem

$$|f(x) - f(0)| < |x - 0|.$$

(If you do not see how to show this, you can still use this result in the next part of the problem.)

- d. What is the basin of attraction of 0?
- e. Show the Schwarzian derivative of f is negative. Note:

$$S_f(x) = \frac{f'''(x)f'(x) - \frac{3}{2}f''(x)^2}{f'(x)^2}.$$

3. (20 Points) Let

$$f(y) = 1 - \frac{3}{4}y^2 \quad \text{and} \quad g(x) = 3x(1-x).$$

Show that $y = C(x) = 4x - 2$ is a conjugacy between f and g . Notice, that x is the variable for g and y is the variable for f .

4. (30 Points) Assume f is a continuous map on the real line which has a period-6 orbit $f(1) = 5$, $f(2) = 6$, $f(3) = 4$, $f(4) = 1$, $f(5) = 2$, and $f(6) = 3$. Label the interval between these points with symbols a through e by $I_a = [1, 2]$, $I_b = [2, 3]$, etc.

a. Give the transition graph for these intervals.

b. What periods are forced to exist by this transition graph? Notice that most orbits alternate between the intervals $[1, 3]$ and $[4, 6]$.

5. (20 Points) Let f be the map

$$f(x) = \frac{4}{\pi} \arctan(x).$$

Note that $f(0) = 0$, $f(1) = 1$, and $f(-1) = -1$. Give all the attracting sets for f . Which of these sets are attractors? Why are there no chaotic attractors? Remember that

$$f'(x) = \frac{4}{\pi(1+x^2)}.$$

6. (50 Points) Let

$$f(x) = \begin{cases} 3 + \frac{3}{2}(x-1) & \text{for } 0 \leq x \leq 1 \\ 3 - 2(x-1) & \text{for } 1 \leq x \leq 2 \\ 1 + \left(\frac{1+\sqrt{5}}{2}\right)(x-2) & \text{for } 2 \leq x. \end{cases}$$

a. Show that $[1, 3]$ has a trapping region.

b. Show that f has a Markov partition on $[1, 3]$. (Don't worry about the image of points outside $[1, 3]$ like $x = 0$ or $x = 4$.)

c. Is f topologically transitive on $[1, 3]$? If so, why?

d. Does f have a chaotic attractor? If so, why?

e. What estimate can you give for the Lyapunov exponents of orbits in $[1, 3]$ which do not pass through the points 1 and 2 where the derivative does not exist? Give an estimate like $A \leq \ell(x_0; f) \leq B$ where A and B are specific values.