

Extended Divergence-Measure Fields and the Euler Equations for Gas Dynamics

Gui-Qiang Chen¹, Hermano Frid²

¹ Department of Mathematics, Northwestern University, 2033 Sheridan Road, Evanston, IL 60208-2730, USA. E-mail: gqchen@math.northwestern.edu

² Instituto de Matemática Pura e Aplicada – IMPA, Estrada Dona Castorina, 110, Rio de Janeiro, RJ 22460-320, Brazil. E-mail: hermano@impa.br

Received: 7 May 2002 / Accepted: 2 December 2002
Published online: 2 April 2003 – © Springer-Verlag 2003

Abstract: A class of extended vector fields, called extended divergence-measure fields, is analyzed. These fields include vector fields in L^p and vector-valued Radon measures, whose divergences are Radon measures. Such extended vector fields naturally arise in the study of the behavior of entropy solutions of the Euler equations for gas dynamics and other nonlinear systems of conservation laws. A new notion of normal traces over Lipschitz deformable surfaces is developed under which a generalized Gauss-Green theorem is established even for these extended fields. An explicit formula is obtained to calculate the normal traces over any Lipschitz deformable surface, suitable for applications, by using the neighborhood information of the fields near the surface and the level set function of the Lipschitz deformation surfaces. As an application, we prove the uniqueness and stability of Riemann solutions that may contain vacuum in the class of entropy solutions of the Euler equations for gas dynamics.

1. Introduction

We are concerned with a class of extended vector fields, called extended divergence-measure fields, or \mathcal{DM} -fields for short. These fields include vector fields in L^p , $1 \leq p \leq \infty$, and vector-valued Radon measures, whose divergences are Radon measures. The \mathcal{DM} -fields naturally arise in the study of the behavior of entropy solutions of nonlinear hyperbolic systems of conservation laws, which take the form

$$u_t + \nabla_x \cdot f(u) = 0, \quad u \in \mathbb{R}^m, \quad x \in \mathbb{R}^n,$$

where $f : \mathbb{R}^m \rightarrow (\mathbb{R}^m)^n$ is a nonlinear map. One of its most important prototypes is the Euler equations for gas dynamics in Lagrangian coordinates:

$$\tau_t - v_x = 0, \tag{1.1}$$

$$v_t + p_x = 0, \tag{1.2}$$

$$\left(e + \frac{v^2}{2} \right)_t + (pv)_x = 0, \tag{1.3}$$

where $\tau = 1/\rho$ is the specific volume with the density ρ , and v, p, e are the velocity, the pressure, the internal energy, respectively; the other two gas dynamical variables are the temperature θ and the entropy S . For ideal polytropic gases, system (1.1)–(1.3) is closed by the following constitutive relations:

$$p\tau = R\theta, \quad e = c_v\theta, \quad p(\tau, S) = \kappa\tau^{-\gamma}e^{S/c_v}, \tag{1.4}$$

where c_v, R , and κ are positive constants, and $\gamma = 1 + c_v/R > 1$. For isentropic gases, the Euler equations become

$$\tau_t - v_x = 0, \tag{1.5}$$

$$v_t + p(\tau)_x = 0, \tag{1.6}$$

where $p(\tau) = \kappa\tau^{-\gamma}, \gamma > 1$.

The main feature of nonlinear hyperbolic conservation laws, especially (1.1)–(1.3), is that, no matter how smooth the initial data is, the solution may develop singularities and form shock waves in finite time. One may expect solutions in the space of functions of bounded variation. This is indeed the case by the Glimm theorem [15] which establishes that, when the initial data has sufficiently small total variation and stays away from vacuum for (1.1)–(1.3), there exists a global entropy solution in BV satisfying the Clausius inequality:

$$S_t \geq 0 \tag{1.7}$$

in the sense of distributions. However, when the initial data is large, still away from vacuum, the solution may develop vacuum in finite time, even instantaneously as $t > 0$. In this case, the specific volume $\tau = 1/\rho$ may then become a Radon measure or an L^1 function, rather than a function of bounded variation. This indicates that solutions of nonlinear hyperbolic conservation laws are generally either in $\mathcal{M}(\mathbb{R}_+ \times \mathbb{R}^n)$, the space of signed Radon measures, or in $L^p(\mathbb{R}_+ \times \mathbb{R}^n), 1 \leq p \leq \infty$. On the other hand, the fact that (1.1)–(1.3) and (1.7) hold in the sense of distributions implies, in particular, that the divergences of the fields $(\tau, -v), (v, p), (e + v^2/2, pv)$, and $(S, 0)$ in the (t, x) variables are Radon measures, in which the first three are the trivial null measure and the last one is a nonnegative measure as a consequence of the Schwartz Lemma [21]. This motivates our study of extended divergence-measure fields (see Definition 1.1 below).

In this connection, we recall that Wagner [25] proved that the well known Lagrangian transformation carries entropy solutions of the Euler equations in Eulerian coordinates to entropy solutions of (1.1)–(1.3) in a one-to-one and onto manner. However, since solutions of the former which contain vacuum are carried into solutions of (1.1)–(1.3) which are vector-valued measures, the concept of entropy solutions for the latter has to be strengthened. We will return to this point in Sect. 4.

Understanding more properties of \mathcal{DM} -fields can advance our understanding of the behavior of solutions (cf. [5–7]). One of the fundamental questions is whether the normal traces can still be defined and the Gauss-Green formula, i.e., integration by parts, still works for these extended fields, which are very weak.

We begin with the definition of \mathcal{DM} -fields. For open sets $A, B \subset \mathbb{R}^N$, by the relation $A \Subset B$ we mean that the closure of A , \bar{A} , is a compact subset of B .

Definition 1.1. Let $D \subset \mathbb{R}^N$ be open. For $F \in L^p(D; \mathbb{R}^N)$, $1 \leq p \leq \infty$, or $F \in \mathcal{M}(D; \mathbb{R}^N)$, set

$$|\operatorname{div} F|(D) := \sup \left\{ \int_D \nabla \varphi \cdot F : \varphi \in C_0^1(D), |\varphi(x)| \leq 1, x \in D \right\}.$$

For $1 \leq p \leq \infty$, we say that F is an L^p -divergence-measure field over D , i.e., $F \in \mathcal{DM}^p(D)$, if $F \in L^p(D; \mathbb{R}^N)$ and

$$\|F\|_{\mathcal{DM}^p(D)} := \|F\|_{L^p(D; \mathbb{R}^N)} + |\operatorname{div} F|(D) < \infty. \tag{1.8}$$

We say that F is an extended divergence-measure field over D , i.e., $F \in \mathcal{DM}^{ext}(D)$, if $F \in \mathcal{M}(D; \mathbb{R}^N)$ and

$$\|F\|_{\mathcal{DM}^{ext}(D)} := |F|(D) + |\operatorname{div} F|(D) < \infty. \tag{1.9}$$

If $F \in \mathcal{DM}^p(D)$ for any open set $D \Subset \mathbb{R}^N$, then we say $F \in \mathcal{DM}_{loc}^p(\mathbb{R}^N)$; and, if $F \in \mathcal{DM}^{ext}(D)$ for any open set $D \Subset \mathbb{R}^N$, we say $F \in \mathcal{DM}_{loc}^{ext}(\mathbb{R}^N)$.

It is easy to check that these spaces under the norms (1.8) and (1.9), respectively, are Banach spaces. These spaces are larger than the BV -space. The establishment of the Gauss–Green theorem, traces, and other properties of BV functions in the middle of last century (see Federer [13]) has advanced significantly our understanding of solutions of nonlinear partial differential equations and nonlinear problems in calculus of variations, differential geometry, and other areas. A natural question is whether the \mathcal{DM} -fields have similar properties, especially the normal traces and the Gauss–Green formula. At a first glance, it seems unclear.

Example 1.1. The field $F(x, y) = \left(\frac{x}{x^2+y^2}, \frac{y}{x^2+y^2} \right)$ belongs to $\mathcal{DM}_{loc}^1(\mathbb{R}^2)$. As remarked in Whitney [26], for $\Omega = (0, 1) \times (0, 1)$,

$$\int_{\Omega} \operatorname{div} F = 0 \neq \int_{\partial\Omega} F \cdot \nu \, d\mathcal{H}^1 = \frac{\pi}{2},$$

if one understands $F \cdot \nu$ in the classical sense, which implies that the classical Gauss–Green theorem fails. In this paper, we succeed in using the neighborhood information via the Lipschitz deformation to develop a natural notion of normal traces, under which our generalized Gauss–Green theorem holds, even for $F \in \mathcal{DM}^{ext}(D)$.

Example 1.2. For any $\mu \in \mathcal{M}(\mathbb{R})$ with finite total variation,

$$F(x, y) = (dx \times \mu(y), 0) \in \mathcal{DM}^{ext}(I \times \mathbb{R}),$$

for any bounded open interval $I \subset \mathbb{R}$. A non-trivial example of such fields is provided by the Riemann solutions of the Euler equations (1.1)–(1.3) for gas dynamics, which develop vacuum. See (4.12) below.

Some efforts have been made in generalizing the Gauss-Green Theorem. Some results for several situations can be found in Anzellotti [1] for an abstract formulation for $F \in L^\infty$, Rodrigues [20] for $F \in L^2$, and Ziemer [28] for a related problem for $\operatorname{div} F \in L^1$ (see also Baiocchi-Capelo [2], Brezzi-Fortin [4], and Ziemer [29]). In Chen-Frid [5], we observed an explicit way to calculate the normal traces for $F \in \mathcal{DM}^\infty$ by the neighborhood information, under which the Gauss-Green formula holds for such fields.

In this paper, motivated by various nonlinear problems from conservation laws, we propose a natural notion of normal traces by the neighborhood information via Lipschitz deformation under which a generalized Gauss-Green theorem is established for $F \in \mathcal{DM}^{ext}(D)$ in Sect. 3, where our main results concerning extended divergence-measure fields are stated and proved, after establishing some auxiliary results in Sect. 2. In particular, we show an explicit way to calculate the normal traces over any deformable Lipschitz surface, suitable for applications, by using the neighborhood information of the fields near the surface and the level set function of the Lipschitz deformation surfaces. We also show a product rule for these extended fields. Their proofs require some refined properties of Radon measures and the Whitney extension theory, among others.

In Sect. 4, we give an important application of the theory of \mathcal{DM} -fields to the Euler equations (1.1)–(1.3) for gas dynamics and establish the uniqueness and stability of Riemann solutions of large oscillation that may contain two rarefaction waves and one contact discontinuity or vacuum states (i.e. measure solutions) in the class of entropy solutions, which may not belong to either BV_{loc} or L^∞ , without specific reference on the method of construction of the solutions. The proof, motivated by [11] and [7–9], is heavily based on our explicit approach to calculate normal traces over Lipschitz deformable surfaces, in the generalized Gauss-Green theorem, and the product rule for \mathcal{DM} -fields. The same arguments clearly also yield the uniqueness and stability of Riemann solutions in the class of entropy solutions for the Euler equations (1.5) and (1.6) for isentropic gas dynamics.

Before closing this introduction, we recall some correlated results. In DiPerna [11], a uniqueness theorem of Riemann solutions was first established for 2×2 systems in the class of entropy solutions in $L^\infty \cap BV_{loc}$ with small oscillation. We also refer to Dafermos [10] for the stability of Lipschitz solutions for hyperbolic systems of conservation laws. In [8, 9], the uniqueness and stability of Riemann solutions of large oscillation without vacuum (possibly containing shocks) was proved for the 3×3 Euler equations, in the class of entropy solutions in $L^\infty \cap BV_{loc}$ which stay away from vacuum. Another related connection is the recent results on the L^1 -stability of the solutions in $L^\infty \cap BV$ obtained either by the Glimm scheme [15], the wave front-tracking method, the vanishing viscosity method, or more generally satisfying an additional regularity, with small total variation in x uniformly for all $t > 0$ (see the recent references cited in Bianchini-Bressan [3] and Dafermos [10]).

2. Radon Measures and the Whitney Extension

In this section, we establish some auxiliary properties about Radon measures and the Whitney extension of Lipschitz continuous functions, which are required for our analysis on extended divergence-measure vector fields in Sect. 3. We begin with some properties about Radon measures. Let $\Omega, D \subset \mathbb{R}^N$ be open. For $\mu^\epsilon, \mu \in \mathcal{M}(\Omega)$, we denote $\mu^\epsilon \rightharpoonup \mu$ the weak convergence of μ^ϵ to μ in $\mathcal{M}(\Omega)$. The next three lemmas are standard, but we include their proofs for completeness.

Lemma 2.1. *Let $\mu^\varepsilon, \mu \in \mathcal{M}(\Omega)$ be signed Radon measures over Ω with $\mu^\varepsilon \rightharpoonup \mu$. Then*

$$|\mu|(\Omega) \leq \liminf_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(\Omega).$$

This can be seen as follows: For any $\phi \in C_0(\Omega)$, $|\phi| \leq 1$,

$$\langle \mu, \phi \rangle = \lim_{\varepsilon \rightarrow 0} \langle \mu^\varepsilon, \phi \rangle \leq \liminf_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(\Omega).$$

Lemma 2.2. *Let $\mu^\varepsilon, \mu \in \mathcal{M}(\Omega)$ be such that*

$$\mu^\varepsilon \rightharpoonup \mu, \quad \lim_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(\Omega) = |\mu|(\Omega).$$

Then, for every open set $A \subset \Omega$,

$$|\mu|(\bar{A} \cap \Omega) \geq \limsup_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(\bar{A} \cap \Omega).$$

In particular, if $|\mu|(\partial A \cap \Omega) = 0$, then

$$|\mu|(A) = \lim_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(A).$$

Proof. Set $B = \Omega - \bar{A}$, which is open. Then

$$|\mu|(A) \leq \liminf_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(A), \quad |\mu|(B) \leq \liminf_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(B).$$

On the other hand,

$$\begin{aligned} |\mu|(\bar{A} \cap \Omega) + |\mu|(B) &= |\mu|(\Omega) = \lim_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(\Omega) \\ &\geq \limsup_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(\bar{A} \cap \Omega) + \liminf_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(B) \\ &\geq \limsup_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(\bar{A} \cap \Omega) + |\mu|(B), \end{aligned}$$

which yields the desired result. \square

Lemma 2.3. *Let ω^ε be a sequence of positive symmetric mollifiers in \mathbb{R}^N and $\mu \in \mathcal{M}(\Omega)$. Set $\mu^\varepsilon = \mu * \omega^\varepsilon$, the mollified measures. Then, for any open set $A \subset \subset \Omega$ with $|\mu|(\partial A) = 0$,*

$$|\mu|(A) = \lim_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(A).$$

Proof. Since $\mu^\varepsilon \rightharpoonup \mu$, Lemma 2.1 implies

$$|\mu|(A) \leq \liminf_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(A).$$

Notice that, for any $g \in C_0(A)$, $|g| \leq 1$,

$$\langle \mu^\varepsilon, g \rangle = \langle \mu, g^\varepsilon \rangle,$$

where $g^\varepsilon = g * \omega^\varepsilon$. Since $|g^\varepsilon| \leq 1$ and $\text{spt}(g^\varepsilon) \subset A_\varepsilon := \{x : \text{dist}(x, A) \leq \varepsilon\} \subset \Omega$ for $\varepsilon \ll 1$, then

$$|\langle \mu^\varepsilon, g \rangle| \leq |\mu|(A_\varepsilon),$$

which implies

$$|\mu^\varepsilon|(A) \leq |\mu|(A_\varepsilon).$$

Hence,

$$\limsup_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(A) \leq \lim_{\varepsilon \rightarrow 0} |\mu|(A_\varepsilon) = |\mu|(\bar{A}),$$

which yields

$$\limsup_{\varepsilon \rightarrow 0} |\mu^\varepsilon|(A) \leq |\mu|(A),$$

since $|\mu|(\partial A) = 0$. \square

In particular, Lemma 2.3 indicates that, if $\mu \geq 0$ and $\mu(\partial A) = 0$, then

$$\mu(A) = \lim_{\varepsilon \rightarrow 0} \mu^\varepsilon(A).$$

Proposition 2.1. *Let $D \subset \mathbb{R}^N$ be open, $\mu \in \mathcal{M}(D)$, and $\mu^\varepsilon = \mu * \omega^\varepsilon$. Let $\Omega \subset\subset D$ be open and $|\mu|(\partial\Omega) = 0$. Then, for any $\phi \in C(D)$,*

$$\lim_{\varepsilon \rightarrow 0} \langle \mu^\varepsilon, \phi \chi_\Omega \rangle = \langle \mu, \phi \chi_\Omega \rangle.$$

Proof. Write

$$\mu^\varepsilon = \mu^{\varepsilon+} - \mu^{\varepsilon-},$$

where $\mu^{\varepsilon\pm} = \mu^\pm * \omega^\varepsilon$ are nonnegative measures. The condition $|\mu|(\partial\Omega) = 0$ implies

$$\mu^\pm(\partial\Omega) = 0.$$

From Lemma 2.3, we have

$$\lim_{\varepsilon \rightarrow 0} \mu^{\varepsilon\pm}(\Omega) = \mu^\pm(\Omega).$$

Hence

$$\lim_{\varepsilon \rightarrow 0} \mu^\varepsilon(\Omega) = \lim_{\varepsilon \rightarrow 0} \mu^{\varepsilon+}(\Omega) - \lim_{\varepsilon \rightarrow 0} \mu^{\varepsilon-}(\Omega) = \mu^+(\Omega) - \mu^-(\Omega) = \mu(\Omega).$$

Let A be open with $\Omega \subset\subset A \subset\subset D$. Then, for $\phi \in C(D)$ and $\delta > 0$, we may construct a partition of \mathbb{R}^N by means of parallelepipeds:

$$Q_\alpha = [x_1^{\alpha_1}, x_1^{\alpha_1+1}] \times \cdots \times [x_N^{\alpha_N}, x_N^{\alpha_N+1}], \quad \alpha = (\alpha_1, \dots, \alpha_N) \in \mathbb{Z}^N,$$

such that

$$\begin{aligned} \cup_{Q_\alpha \cap \Omega \neq \emptyset} Q_\alpha &\subset A, \\ |\mu|(\partial Q_\alpha \cap D) &= 0, \quad \alpha \in \mathbb{Z}^N, \end{aligned}$$

and

$$|\phi(x) - \phi(y)| < \delta, \quad x, y \in Q_\alpha.$$

Let $a_\alpha = \phi(x_\alpha^*)$ for some $x_\alpha^* \in Q_\alpha$ if $Q_\alpha \cap \Omega \neq \emptyset$, and $a_\alpha = 0$, otherwise. Then

$$\langle \mu^\varepsilon, \phi \chi_\Omega \rangle = \left\langle \mu^\varepsilon, \sum a_\alpha \chi_{Q_\alpha \cap \Omega} \right\rangle + \left\langle \mu^\varepsilon, \left(\phi - \sum a_\alpha \chi_{Q_\alpha} \right) \chi_\Omega \right\rangle,$$

and

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} \langle \mu^\varepsilon, \phi \chi_\Omega \rangle &\leq \left\langle \mu, \sum a_\alpha \chi_{Q_\alpha \cap \Omega} \right\rangle + \delta |\mu|(A) \\ &\leq \langle \mu, \phi \chi_\Omega \rangle + 2\delta |\mu|(A). \end{aligned}$$

Analogously,

$$\liminf_{\varepsilon \rightarrow 0} \langle \mu^\varepsilon, \phi \chi_\Omega \rangle \geq \langle \mu, \phi \chi_\Omega \rangle - 2\delta |\mu|(A).$$

Since $\delta > 0$ is arbitrary, we complete the proof. \square

We now discuss some concepts and facts about the extension of Lipschitz continuous functions defined on a closed set $C \subset \mathbb{R}^N$ to \mathbb{R}^N , following the theory set forth by Whitney [27] (see also [23]), which play an important role in Sect. 3.

Let k be a nonnegative integer and $\gamma \in (k, k + 1]$. We say that a function f , defined on C , belongs to $\text{Lip}(\gamma, C)$ if there exist functions $f^{(j)}$, $0 \leq |j| \leq k$, defined on C , with $f^{(0)} = f$, such that, if

$$f^{(j)}(x) = \sum_{|j+l| \leq k} \frac{f^{(j+l)}(y)}{l!} (x - y)^l + R_j(x, y),$$

then

$$\begin{cases} |f^{(j)}(x)| \leq M, \\ |R_j(x, y)| \leq M|x - y|^{\gamma - |j|}, \quad \text{for any } x, y \in C, |j| \leq k. \end{cases} \tag{2.1}$$

Here j and l denote multi-indices $j = (j_1, \dots, j_N)$ and $l = (l_1, \dots, l_N)$ with $j! = j_1! \cdots j_N!$, $|j| = j_1 + j_2 + \dots + j_N$, and $x^l = x_1^{l_1} x_2^{l_2} \cdots x_N^{l_N}$. By an element of $\text{Lip}(\gamma, C)$ we mean the collection $\{f^{(j)}(x)\}_{|j| \leq k}$. The norm of an element in $\text{Lip}(\gamma, C)$ is defined as the smallest M for which inequality (2.1) holds. We notice that $\text{Lip}(\gamma, C)$ with this norm is a Banach space. For the case $C = \mathbb{R}^N$, since the functions $f^{(j)}$ are determined by $f^{(0)}$, this collection is then identified with $f^{(0)}$.

The Whitney extension of order k is defined as follows. Let $\{f^{(j)}\}_{|j| \leq k}$ be an element of $\text{Lip}(\gamma, C)$. The linear mapping $\mathcal{E}_k : \text{Lip}(\gamma, C) \rightarrow \text{Lip}(\gamma, \mathbb{R}^N)$ assigns to any such collection a function $\mathcal{E}_k(f^{(j)})$ defined on \mathbb{R}^N which is an extension of $f^{(0)} = f$ to \mathbb{R}^N . The definition of \mathcal{E}_k is the following:

$$\begin{cases} \mathcal{E}_0(f)(x) = f(x) & \text{for } x \in C, \\ \mathcal{E}_0(f)(x) = \sum_i f(p_i) \varphi_i(x) & \text{for } x \in \mathbb{R}^N - C, \end{cases}$$

and, for $k > 0$,

$$\begin{cases} \mathcal{E}_k(f^{(j)})(x) = f^{(0)}(x) & \text{for } x \in C, \\ \mathcal{E}_k(f^{(j)})(x) = \sum_i P(x, p_i) \varphi_i(x) & \text{for } x \in \mathbb{R}^N - C. \end{cases}$$

Here $P(x, y)$ denotes the polynomial in x , which is the Taylor expansion of f about the point $y \in C$:

$$P(x, y) = \sum_{|l| \leq k} \frac{f^{(l)}(y)(x - y)^l}{l!} \quad \text{for } x \in \mathbb{R}^N, y \in C.$$

The functions φ_i form a partition of unity of $\mathbb{R}^N - C$ with the following properties:

(i) $\text{spt}(\varphi_i) \subset Q_i$, where Q_i is a cube with edges parallel to the coordinate axes and

$$c_1 \text{diam}(Q_i) \leq \text{dist}(Q_i, C) \leq c_2 \text{diam}(Q_i),$$

for certain positive constants c_1 and c_2 independent of C ;

- (ii) each point of $\mathbb{R}^N - C$ is contained in at most N_0 cubes Q_i , for a certain number N_0 depending only on the dimension N ;
- (iii) the derivatives of φ_i satisfy

$$|\partial_{x_1}^{\alpha_1} \cdots \partial_{x_N}^{\alpha_N} \varphi_i(x)| \leq A_\alpha (\text{diam } Q_i)^{-|\alpha|} \quad \text{for } x \in Q_i. \tag{2.2}$$

Here $p_i \in C$ is such that $\text{dist}(Q_i, C) = \text{dist}(p_i, Q_i)$, $|\alpha| = \alpha_1 + \cdots + \alpha_N$, and the symbol \sum' indicates that the summation is taken only over those cubes whose distances to C are not greater than one.

The following theorem, whose proof can also be found in [23], is due to Whitney [27].

Theorem 2.1. *Suppose that k is a nonnegative integer, $\gamma \in (k, k + 1]$, and C is a closed set. Then the mapping \mathcal{E}_k is a continuous linear mapping from $\text{Lip}(\gamma, C)$ to $\text{Lip}(\gamma, \mathbb{R}^N)$ which defines an extension of $f^{(0)}$ to \mathbb{R}^N , and the norm of this mapping has a bound independent of C .*

We will need the following proposition, which seems to be of interest in itself and is useful in establishing the generalized Gauss-Green theorem in Sect. 3.

Proposition 2.2. *Let C be a closed set in \mathbb{R}^N and*

$$C_\delta := \{x \in \mathbb{R}^N : \text{dist}(x, C) \leq \delta\} \quad \text{for } \delta > 0.$$

Let $\mathcal{E}_k : \text{Lip}(\gamma, C) \rightarrow \text{Lip}(\gamma, \mathbb{R}^N)$ with $\gamma \in (k, k + 1]$ be the Whitney extension of order k . Then, for any $\phi \in \text{Lip}(\gamma, \mathbb{R}^N)$ and any $\gamma' \in (k, \gamma)$,

$$\|\mathcal{E}_k(\phi|C) - \phi\|_{\text{Lip}(\gamma', C_\delta)} \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \tag{2.3}$$

Proof. We will prove the proposition in detail only for the cases $k = 0$ and $k = 1$ since the case $k > 1$ can then be obtained by induction.

For $k = 0$, \mathcal{E}_0 is given by $\mathcal{E}_0(f)(x) = f(x)$ if $x \in C$, and

$$\mathcal{E}_0(f)(x) = \sum_{i=1}^{\infty} f(p_i)\varphi_i(x) \quad \text{if } x \in \mathbb{R}^N - C, \tag{2.4}$$

where p_i and φ_i are as above. Now, for any $\phi \in \text{Lip}(\gamma, \mathbb{R}^N)$, we have

$$\mathcal{E}_0(\phi|C)(x) - \phi(x) = \sum_{i=1}^{\infty} (\phi(p_i) - \phi(x))\varphi_i(x).$$

Clearly,

$$\sup_{x \in C_\delta} |\mathcal{E}_0(\phi|C)(x) - \phi(x)| \leq c \sup_{\substack{y \in C, x \in C_\delta \\ |x-y| \leq c_0\delta}} |\phi(y) - \phi(x)| \rightarrow 0, \quad \text{as } \delta \rightarrow 0. \tag{2.5}$$

Here and in what follows in this proof c_0 and c are positive constants which are independent of $\delta > 0$ and C , whose values may change at each appearance.

Set $g = \mathcal{E}_0(\phi|C) - \phi$. We now show that

$$|g(x) - g(y)| \leq M(\delta)|x - y|^{\gamma'} \quad \text{for } x, y \in C_\delta, \tag{2.6}$$

where $M(\delta) \rightarrow 0$ as $\delta \rightarrow 0$. Indeed,

$$g(x) - g(y) = \sum_{x \in Q_i} (\phi(p_i) - \phi(x))\varphi_i(x) - \sum_{y \in Q_i} (\phi(p_i) - \phi(y))\varphi_i(y).$$

We split each of these sums into two, respectively:

$$\sum_{x \in Q_i} = \sum_{\substack{x \in Q_i \\ y \notin Q_i}} + \sum_{\substack{x \in Q_i \\ y \in Q_i}}, \quad \sum_{y \in Q_i} = \sum_{\substack{y \in Q_i \\ x \notin Q_i}} + \sum_{\substack{y \in Q_i \\ x \in Q_i}},$$

and denote

$$\sum^0 = \sum_{\substack{x \in Q_i \\ y \in Q_i}}, \quad \sum^1 = \sum_{\substack{x \in Q_i \\ y \notin Q_i}}, \quad \sum^2 = \sum_{\substack{y \in Q_i \\ x \notin Q_i}}.$$

We have

$$\begin{aligned} & \sum^0 ((\phi(p_i) - \phi(x))\varphi_i(x) - (\phi(p_i) - \phi(y))\varphi_i(y)) \\ &= \sum^0 (\phi(y) - \phi(x))\varphi_i(x) + \sum^0 (\phi(p_i) - \phi(y))(\varphi_i(x) - \varphi_i(y)). \end{aligned}$$

Since, if $x, y \in Q_i \cap C_\delta$, $|x - y| \leq c_0\delta$, for a given constant $c_0 > 0$, we obtain, for the first sum,

$$\left| \sum^0 (\phi(x) - \phi(y))\varphi_i(x) \right| \leq c\delta^{\gamma-\gamma'}|x - y|^{\gamma'},$$

and, for the second sum,

$$\begin{aligned} \left| \sum^0 (\phi(p_i) - \phi(y))(\varphi_i(x) - \varphi_i(y)) \right| &\leq c \sum^0 |p_i - y|^\gamma (\text{diam } Q_i)^{-1} |x - y| \\ &\leq c |x - y|^\gamma \leq c \delta^{\gamma-\gamma'} |x - y|^{\gamma'}. \end{aligned}$$

Now, for \sum^1 , we have

$$\sum^1 (\phi(p_i) - \phi(x))\varphi_i(x) = \sum^1 (\phi(p_i) - \phi(x))(\varphi_i(x) - \varphi_i(q_i)),$$

with $q_i = \partial Q_i \cap [x, y]$, where $[x, y]$ is the straight line segment connecting x to y . Therefore,

$$\begin{aligned} \sum^1 |\phi(p_i) - \phi(x)| |\varphi_i(x) - \varphi_i(q_i)| &\leq c \sum^1 |p_i - x|^\gamma (\text{diam } Q_i)^{-1} |x - q_i| \\ &\leq c \sum^1 |x - q_i|^\gamma \leq c \delta^{\gamma-\gamma'} |x - y|^{\gamma'}. \end{aligned}$$

The sum \sum^2 is treated similarly. We have then proved (2.6) which, together with (2.5), gives (2.3) for $k = 0$.

For $k = 1$, we have $\mathcal{E}_1(f)(x) = f(x)$ for $x \in C$, and

$$\mathcal{E}_1(f)(x) = \sum_{i=1}^{\infty} P_f(x, p_i)\varphi_i(x) \quad \text{for } x \in \mathbb{R}^N - C,$$

where

$$P_f(x, y) = f(y) + \sum_{j=1}^N \partial_{x_j} f(y)(x_j - y_j),$$

and the functions φ_i , conveniently renumbered together with Q_i containing $\text{spt}(\varphi_i)$, satisfy $\text{dist}(Q_i, C) \leq 1$. Since $\phi \in \text{Lip}(\gamma, \mathbb{R}^N)$, $\gamma > 1$, we clearly have

$$(\phi|C)^j(y) = \partial_{x_j} \phi(y) \quad \text{for } y \in C.$$

Hence

$$\mathcal{E}_1(\phi|C)(x) - \phi(x) = \sum_{i=1}^{\infty} (P_{\phi}(x, p_i) - \phi(x))\varphi_i(x).$$

Setting $g_1 = \mathcal{E}_1(\phi|C) - \phi$, we have

$$\begin{aligned} |g_1(x)| &\leq \sum_{i=1}^{\infty} |P_{\phi}(x, p_i) - \phi(x)| |\varphi_i(x)| \\ &\leq c \sum_{x \in Q_i} |x - p_i|^{\gamma} |\varphi_i(x)| \leq c \delta^{\gamma} \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \end{aligned}$$

Also,

$$\begin{aligned} \partial_{x_k} g_1(x) &= \sum_{x \in Q_i} \partial_{x_k} (P_{\phi}(x, p_i) - \phi(x))\varphi_i(x) + \sum_{x \in Q_i} (P_{\phi}(x, p_i) - \phi(x))\partial_{x_k} \varphi_i(x) \\ &= h_1(x) + h_2(x), \end{aligned}$$

where

$$h_1(x) := \sum_{x \in Q_i} (\partial_{x_k} \phi(p_i) - \partial_{x_k} \phi(x))\varphi_i(x), \quad h_2(x) := \sum_{x \in Q_i} (P_{\phi}(x, p_i) - \phi(x))\partial_{x_k} \varphi_i(x).$$

Hence,

$$\begin{aligned} |\partial_{x_k} g_1(x)| &\leq c \sum_{x \in Q_i} |p_i - x|^{\gamma-1} + c \sum_{x \in Q_i} |x - p_i|^{\gamma} \text{diam}(Q_i)^{-1} \\ &\leq c \delta^{\gamma-1} \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \end{aligned}$$

We now show

$$|\partial_{x_k} g_1(x) - \partial_{x_k} g_1(y)| \leq M(\delta)|x - y|^{\gamma'-1},$$

with $M(\delta) \rightarrow 0$ as $\delta \rightarrow 0$. First we obtain

$$|h_1(x) - h_1(y)| \leq M(\delta)|x - y|^{\gamma'-1}, \quad \text{with } M(\delta) \rightarrow 0 \quad \text{as } \delta \rightarrow 0,$$

exactly as in the case $k = 0$. Now,

$$\begin{aligned} h_2(x) - h_2(y) &= \sum_{x \in Q_i} (P_\phi(x, p_i) - \phi(x)) \partial_{x_k} \phi_i(x) - \sum_{x \in Q_i} (P_\phi(x, p_i) - \phi(y)) \partial_{x_k} \phi_i(y) \\ &= \sum_{x \in Q_i} R(x, p_i) \partial_{x_k} \phi_i(x) - \sum_{y \in Q_i} R(y, p_i) \partial_{x_k} \phi_i(y), \end{aligned}$$

where we set $R(x, y) := P_\phi(x, y) - \phi(x)$. Again, we split each of the last two sums above into two, respectively:

$$\sum_{x \in Q_i} = \sum^0 + \sum^1, \quad \sum_{y \in Q_i} = \sum^0 + \sum^2,$$

as in the first part of the proof, and compute

$$\begin{aligned} & \left| \sum^0 R(x, p_i) \partial_{x_k} \phi_i(x) - \sum^0 R(y, p_i) \partial_{x_k} \phi_i(y) \right| \\ & \leq \left| \sum^0 (R(x, p_i) - R(y, p_i)) \partial_{x_k} \phi_i(x) \right| + \left| \sum^0 R(y, p_i) (\partial_{x_k} \phi_i(y) - \partial_{x_k} \phi_i(x)) \right| \\ & \leq c \sum^0 |x - y|^\gamma (\text{diam } Q_i)^{-1} + c \sum^0 |y - p_i|^\gamma (\text{diam } Q_i)^{-2} |x - y| \\ & \leq c \delta^{\gamma - \gamma'} |x - y|^{\gamma' - 1}. \end{aligned}$$

For the remaining sums, we have

$$\begin{aligned} \left| \sum^1 R(x, p_i) \partial_{x_k} \phi_i(x) \right| &= \left| \sum^1 R(x, p_i) (\partial_{x_k} \phi_i(x) - \partial_{x_k} \phi_i(q_i)) \right| \\ &\leq c \sum^1 |x - p_i|^\gamma (\text{diam } Q_i)^{-2} |x - q_i| \\ &\leq c \delta^{\gamma - \gamma'} |x - y|^{\gamma' - 1}, \end{aligned}$$

where q_i is as that in the first part of the proof; and the sum $\sum^2 R(y, p_i) \partial_{x_k} \phi_i(y)$ is treated similarly. This concludes the proof in the case $k = 1$. As indicated above, the case $k > 1$ follows similarly by induction. \square

3. Normal Traces and Generalized Gauss-Green Theorem

In this section, we prove our main results concerning extended \mathcal{DM} -fields, including a new notion of normal traces, a generalized Gauss-Green theorem, and a product rule for \mathcal{DM} -fields. We begin with the definition of deformable Lipschitz boundaries.

Definition 3.1. *Let $\Omega \subset \mathbb{R}^N$ be an open bounded subset. We say that $\partial\Omega$ is a deformable Lipschitz boundary, provided that*

(i) $\forall x \in \partial\Omega, \exists r > 0$ and a Lipschitz map $\gamma : \mathbb{R}^{N-1} \rightarrow \mathbb{R}$ such that, after rotating and relabeling coordinates if necessary,

$$\Omega \cap Q(x, r) = \{y \in \mathbb{R}^N : \gamma(y_1, \dots, y_{N-1}) < y_N\} \cap Q(x, r),$$

where $Q(x, r) = \{y \in \mathbb{R}^N : |x_i - y_i| \leq r, i = 1, \dots, N\}$;

(ii) $\exists \Psi : \partial\Omega \times [0, 1] \rightarrow \bar{\Omega}$ such that Ψ is a homeomorphism bi-Lipschitz over its image and $\Psi(\omega, 0) = \omega$ for all $\omega \in \partial\Omega$. The map Ψ is called a Lipschitz deformation of the boundary $\partial\Omega$.

The following lemma is a direct corollary of the boundedness of F and $\operatorname{div} F$ over Ω as Radon measures. Since the theory of \mathcal{DM}^∞ -fields has been addressed in [5], henceforth we focus on \mathcal{DM}^* -fields, where $*$ stands for either $p \in [1, \infty)$ or *ext*.

Lemma 3.1. *Let $F \in \mathcal{DM}^*(\Omega)$ with Ω an open set whose boundary $\partial\Omega$ has a Lipschitz deformation Ψ with $\partial\Omega_s = \Psi_s(\partial\Omega)$, $s \in [0, 1]$. Then there exists a countable set $T \subset (0, 1)$ such that*

$$|F|(\partial\Omega_s) = |\operatorname{div} F|(\partial\Omega_s) = 0 \quad \text{for any } s \in (0, 1) - T.$$

We now establish the generalized Gauss-Green theorem for \mathcal{DM}^* -fields, by introducing a suitable definition of normal traces over the boundary $\partial\Omega$ of a bounded open set Ω with Lipschitz deformable boundary.

Theorem 3.1 (Normal Traces and Generalized Gauss-Green Theorem). *Assume $F \in \mathcal{DM}^*(\Omega)$. Let $\Omega \subset \mathbb{R}^N$ is a bounded open set with Lipschitz deformable boundary. Then there exists a continuous linear functional $F \cdot \nu|_{\partial\Omega}$ over $\operatorname{Lip}(\gamma, \partial\Omega)$, $\gamma > 1$, such that, for any $\phi \in \operatorname{Lip}(\gamma, \mathbb{R}^N)$,*

$$\langle F \cdot \nu|_{\partial\Omega}, \phi \rangle = \int_{\Omega} \phi \operatorname{div} F + \int_{\Omega} \nabla \phi \cdot F. \tag{3.1}$$

Let $h : \mathbb{R}^N \rightarrow \mathbb{R}$ be the level set function of $\partial\Omega_s$, that is,

$$h(x) := \begin{cases} 0, & \text{for } x \in \mathbb{R}^N - \overline{\Omega}, \\ 1, & \text{for } x \in \Omega - \Psi(\partial\Omega \times [0, 1]), \\ s, & \text{for } x \in \partial\Omega_s, 0 \leq s \leq 1. \end{cases}$$

If $F \in \mathcal{DM}^p(\Omega)$, $1 \leq p < \infty$, then

$$\begin{aligned} \langle F \cdot \nu|_{\partial\Omega}, \psi \rangle &= - \lim_{s \rightarrow 0} \frac{1}{s} \int_{\Psi(\partial\Omega \times (0, s))} \mathcal{E}(\psi) \nabla h \cdot F \, dx \\ &= - \lim_{s \rightarrow 0} \frac{1}{s} \int_{\Psi(\partial\Omega \times (0, s))} \mathcal{E}(\psi) |\nabla h| F \cdot \nu \, dx \end{aligned} \tag{3.2}$$

for any $\psi \in \operatorname{Lip}(\partial\Omega)$, where $\mathcal{E}(\psi)$ is any Lipschitz extension of ψ to all \mathbb{R}^N and $\nu : \Psi(\partial\Omega \times [0, 1]) \rightarrow \mathbb{R}^N$ is such that $\nu(x)$ is the unit outer normal vector to $\partial\Omega_s$ at $x \in \partial\Omega_s$, defined for a.e. $x \in \Psi(\partial\Omega \times [0, 1])$. Formula (3.2) also holds if $F \in \mathcal{DM}^{ext}(\Omega)$, for any $\psi \in \operatorname{Lip}(\gamma, \partial\Omega)$, $\gamma > 1$, and $\mathcal{E}(\psi) \in \operatorname{Lip}(\gamma, \mathbb{R}^N)$ any extension of ψ to \mathbb{R}^N , provided that the set of non-Lebesgue points of $\nabla h(x)$ on $\Psi(\partial\Omega \times (0, 1))$ has $|F|$ -measure zero. Finally, for $F \in \mathcal{DM}^p(\Omega)$ with $1 < p < \infty$, $F \cdot \nu|_{\partial\Omega}$ can be extended to a continuous linear functional over $W^{1-1/p, p}(\partial\Omega) \cap C(\partial\Omega)$.

Proof. We divide the proof into four steps.

Step 1. We first treat the more general case $F \in \mathcal{DM}^{ext}(\Omega)$. For $\psi \in \operatorname{Lip}(\gamma, \partial\Omega)$, let $\mathcal{E}(\psi)$ be the Whitney extension of ψ of order $k = 1$. Then, by the classical Gauss-Green formula, we have

$$\int_{\partial\Omega} F^\varepsilon \cdot \nu \, \psi \, d\mathcal{H}^{N-1} = \int_{\Omega} \mathcal{E}(\psi) \operatorname{div} F^\varepsilon + \int_{\Omega} \nabla \mathcal{E}(\psi) \cdot F^\varepsilon, \tag{3.3}$$

where $F^\varepsilon = F * \omega_\varepsilon$ with the standard mollified sequence ω_ε .

To begin with, we first focus on $F \in \mathcal{DM}^{ext}(D)$, for $\Omega \Subset D \Subset \mathbb{R}^N$, satisfying

$$|F|(\partial\Omega) = |\operatorname{div} F|(\partial\Omega) = 0. \tag{3.4}$$

The right-hand side of (3.3) defines a uniformly bounded family of continuous linear operators l^ε over $\operatorname{Lip}(\gamma, \partial\Omega)$. Moreover, for each $\psi \in \operatorname{Lip}(\gamma, \partial\Omega)$, the limit $\lim_{\varepsilon \rightarrow 0} l^\varepsilon(\psi)$ exists and equals

$$\int_{\Omega} \mathcal{E}(\psi) \operatorname{div} F + \int_{\Omega} \nabla \mathcal{E}(\psi) \cdot F,$$

as a consequence of (3.4) and Proposition 2.1. Hence, we may define

$$\langle F \cdot \nu|_{\partial\Omega}, \psi \rangle = \lim_{\varepsilon \rightarrow 0} l^\varepsilon(\psi) \quad \text{for any } \psi \in \operatorname{Lip}(\gamma, \partial\Omega).$$

Now, for any $\phi \in \operatorname{Lip}(\gamma, \mathbb{R}^N)$, we let $\varepsilon \rightarrow 0$ in the Gauss-Green formula:

$$\int_{\partial\Omega} F^\varepsilon \cdot \nu \phi \, d\mathcal{H}^{N-1} = \int_{\Omega} \phi \operatorname{div} F^\varepsilon + \int_{\Omega} \nabla \phi \cdot F^\varepsilon,$$

and obtain the identity (3.1).

For the general case that $F \in \mathcal{DM}^{ext}(\Omega)$ without the assumption (3.4), we consider a Lipschitz deformation of $\partial\Omega$, $\Psi : \partial\Omega \times [0, 1] \rightarrow \overline{\Omega}$. Let $s \in (0, 1)$ be such that

$$|F|(\partial\Omega_s) = |\operatorname{div} F|(\partial\Omega_s) = 0. \tag{3.5}$$

Since we have $\mathcal{H}^{N-1}(\partial\Omega_s) < +\infty$, Federer’s extension of the Gauss-Green theorem (see [13]) holds for ϕF^ε over Ω_s . Thus, we know from the previous analysis that $F \cdot \nu|_{\partial\Omega_s}$ is defined as a continuous linear functional over $\operatorname{Lip}(\gamma, \partial\Omega_s)$, whose norm is bounded, independent of $s \in (0, 1)$. Now, for $\psi \in \operatorname{Lip}(\gamma, \partial\Omega)$, we have

$$\langle F \cdot \nu|_{\partial\Omega_s}, (\mathcal{E}(\psi)|_{\partial\Omega_s}) \rangle = \int_{\Omega_s} \mathcal{E}(\psi) \operatorname{div} F + \int_{\Omega_s} \nabla \mathcal{E}(\psi) \cdot F. \tag{3.6}$$

Again, the right-hand side of (3.6) defines a uniformly bounded family of continuous linear functionals l^s over $\operatorname{Lip}(\gamma, \partial\Omega)$, for $s \in (0, 1) - \mathcal{T}$, where \mathcal{T} is defined in Lemma 3.1. Furthermore, $\lim_{s \rightarrow 0} l^s(\psi)$ exists for any $\psi \in \operatorname{Lip}(\gamma, \partial\Omega)$, as a consequence of the Dominated Convergence Theorem applied to both integrals on the right-hand side of (3.6). Hence, we may define

$$\langle F \cdot \nu|_{\partial\Omega}, \psi \rangle = \lim_{s \rightarrow 0} l^s(\psi),$$

which is then a continuous linear functional over $\operatorname{Lip}(\gamma, \partial\Omega)$. Finally, for any $\phi \in \operatorname{Lip}(\gamma, \mathbb{R}^N)$, we obtain (3.1) by taking the limit as $s \rightarrow 0$ in the formula:

$$\langle F \cdot \nu|_{\partial\Omega_s}, (\phi|_{\partial\Omega_s}) \rangle = \int_{\Omega_s} \phi \operatorname{div} F + \int_{\Omega_s} \nabla \phi \cdot F,$$

observing that

$$\begin{aligned} & | \langle F \cdot \nu|_{\partial\Omega_s}, (\phi|_{\partial\Omega_s}) - (\mathcal{E}(\phi|_{\partial\Omega})|_{\partial\Omega_s}) \rangle | \\ & \leq c \| (\phi|_{\partial\Omega_s}) - (\mathcal{E}(\phi|_{\partial\Omega})|_{\partial\Omega_s}) \|_{\operatorname{Lip}(\gamma', \partial\Omega_s)} \\ & \leq c \| \phi - \mathcal{E}(\phi|_{\partial\Omega}) \|_{\operatorname{Lip}(\gamma', \Psi(\partial\Omega \times [0, s]))} \rightarrow 0, \quad \text{as } s \rightarrow 0, \end{aligned}$$

for $1 < \gamma' < \gamma$, as a consequence of Proposition 2.2.

Step 2. We now consider the more regular case that $F \in \mathcal{DM}^p(\Omega)$, $1 \leq p < \infty$. Let F^ε be as above. Again, for any $\phi \in \text{Lip}(\mathbb{R}^N)$, we have

$$\int_{\partial\Omega_s} \phi F^\varepsilon \cdot \nu \, d\mathcal{H}^{N-1} = \int_{\Omega_s} \phi \operatorname{div} F^\varepsilon \, dx + \int_{\Omega_s} \nabla\phi \cdot F^\varepsilon \, dx. \tag{3.7}$$

Now we integrate (3.7) in $s \in (0, \delta)$, $0 < \delta < 1$, and use the coarea formula (see, e.g. [12, 13]) in the left-hand side to obtain

$$-\int_{\Psi(\partial\Omega \times (0, \delta))} \phi F^\varepsilon \cdot \nabla h \, dx = \int_0^\delta \left\{ \int_{\Omega_s} \phi \operatorname{div} F^\varepsilon \, dx \right\} ds + \int_0^\delta \left\{ \int_{\Omega_s} \nabla\phi \cdot F^\varepsilon \, dx \right\} ds. \tag{3.8}$$

Let $\varepsilon \rightarrow 0$. Observing that, by Proposition 2.1, the integrand of the first integral converges for a.e. $s \in (0, \delta)$ to the corresponding integral for F , we obtain

$$-\int_{\Psi(\partial\Omega \times (0, \delta))} \phi F \cdot \nabla h \, dx = \int_0^\delta \left\{ \int_{\Omega_s} \phi \operatorname{div} F \right\} ds + \int_0^\delta \left\{ \int_{\Omega_s} \nabla\phi \cdot F \, dx \right\} ds. \tag{3.9}$$

We then divide (3.9) by δ , let $\delta \rightarrow 0$, and observe that both terms in the right-hand side converge to the corresponding integrals inside the brackets over Ω , by the dominated convergence theorem. Hence, the left-hand side also converges, which yields

$$-\lim_{\delta \rightarrow 0} \frac{1}{\delta} \int_{\Psi(\partial\Omega \times (0, \delta))} \phi F \cdot \nabla h \, dx = \int_{\Omega} \phi \operatorname{div} F + \int_{\Omega} \nabla\phi \cdot F \, dx. \tag{3.10}$$

Now, for $\psi \in \text{Lip}(\partial\Omega)$, let $\mathcal{E}(\psi) \in \text{Lip}(\mathbb{R}^N)$ be a Lipschitz extension of ψ preserving the norm $\|\cdot\|_{\text{Lip}} := \|\cdot\|_\infty + \text{Lip}(\cdot)$ (see, e.g., [12, 13]). We then define

$$\langle F \cdot \nu|_{\partial\Omega}, \psi \rangle = -\lim_{s \rightarrow 0} \frac{1}{s} \int_{\Psi(\partial\Omega \times (0, s))} \mathcal{E}(\psi) \nabla h \cdot F \, dx. \tag{3.11}$$

Because the right-hand side of (3.10) is independent of the particular deformation Ψ over $\partial\Omega$, we see that the normal trace defined by (3.11) is also independent of the deformation.

We still have to prove that the normal trace as defined by (3.11) is also independent of the specific Lipschitz extension $\mathcal{E}(\psi)$ of ψ . This will be accomplished if we prove that the right-hand side of (3.10) vanishes for $\phi|_{\partial\Omega} \equiv 0$. Denote it by $[F, \phi]_{\partial\Omega}$, that is,

$$[F, \phi]_{\partial\Omega} := \langle \operatorname{div} F, \phi \rangle_\Omega + \langle F, \nabla\phi \rangle_\Omega.$$

We claim that $[F, \phi]_{\partial\Omega} = 0$ if $\phi|_{\partial\Omega} \equiv 0$. In fact, we may approximate such ϕ by a sequence $\phi^j \in C_0^\infty(\Omega)$, with $\|\phi^j\|_\infty \leq \|\phi\|_\infty$, such that $\phi^j \rightarrow \phi$ locally uniformly in Ω and $\nabla\phi^j \rightarrow \nabla\phi$ in $L^q(\Omega)^N$, for $p > 1$, $\nabla\phi^j \overset{*}{\rightharpoonup} \nabla\phi$ in $L^\infty(\Omega)^N$, for $p = 1$, with $\frac{1}{p} + \frac{1}{q} = 1$. Hence, $[F, \phi]_{\partial\Omega} = \lim_{j \rightarrow \infty} [F, \phi^j] = 0$, as asserted. In particular, for $1 \leq p \leq \infty$, the values of the normal trace, $\langle F \cdot \nu|_{\partial\Omega}, \phi|_{\partial\Omega} \rangle$, depend only on the values of ϕ over $\partial\Omega$.

Step 3. The fact that formula (3.2) also holds if $F \in \mathcal{DM}^{ext}(\Omega)$, for any $\psi \in \text{Lip}(\gamma, \partial\Omega)$, $\gamma > 1$, provided that the set of non-Lebesgue points of $\nabla h(x)$ on $\Psi(\partial\Omega \times (0, 1))$ has $|F|$ -measure zero, is clear from the above proof for \mathcal{DM}^p -fields, $1 \leq p \leq \infty$.

Step 4. As for the last assertion, we recall a well-known result of Gagliardo [14] which indicates, in particular, that, if $\partial\Omega$ is Lipschitz (that is, satisfies (i) of Definition 3.1) and $\psi \in W^{1-1/p, p}(\partial\Omega)$, then it can be extended into Ω to a function $\mathcal{E}(\psi) \in W^{1, p}(\Omega)$, and

$$\|\mathcal{E}(\psi)\|_{W^{1, p}(\Omega)} \leq c \|\psi\|_{W^{1-1/p, p}(\partial\Omega)}, \tag{3.12}$$

for some positive constant c independent of ψ . From the definition of $\mathcal{E}(\psi)$ given in [14], it is easy to verify that, when $\psi \in C(\partial\Omega)$, $\mathcal{E}(\psi) \in C(\Omega)$ and $\|\mathcal{E}(\psi)\|_{L^\infty(\Omega)} \leq \|\psi\|_{L^\infty(\partial\Omega)}$. Hence, using these facts and (3.10), we easily deduce the last assertion. \square

Remark 3.1. When $F \in \mathcal{DM}^\infty(D)$, the normal trace $F \cdot \nu|_{\partial\Omega}$ is in fact in $L^\infty(\partial\Omega, \mathcal{H}^{N-1})$ as the weak-star limit of $(F \cdot \nu_s) \circ \Psi_s$ in $L^\infty(\partial\Omega, \mathcal{H}^{N-1})$ for any Lipschitz deformation Ψ_s :

$$F \cdot \nu|_{\partial\Omega} = w^* - \lim (F \cdot \nu_s) \circ \Psi_s, \quad L^\infty(\partial\Omega, \mathcal{H}^{N-1}),$$

which is independent of Ψ_s ; and the weak-star topology for this limit is optimal to define $F \cdot \nu|_{\partial\Omega}$ in general (see Chen-Frid [5]). However, for $F \in \mathcal{DM}^*(D)$, the normal traces $F \cdot \nu|_{\partial\Omega}$ may no longer be functions in general. This can be seen in Example 1.1 for $F \in \mathcal{DM}_{loc}^1(\mathbb{R}^2)$ with $\Omega = (0, 1) \times (0, 1)$, for which

$$F \cdot \nu|_{\partial\Omega} = \frac{\pi}{2} \delta_{(0,0)} - d(\mathcal{H}^1 \llcorner \partial\Omega),$$

where $\mathcal{H}^1 \llcorner \partial\Omega$ is the one-dimensional Hausdorff measure on $\partial\Omega$.

Remark 3.2. As mentioned in the proof of Theorem 3.1, if $F \in \mathcal{DM}^p(\Omega)$, for $1 \leq p \leq \infty$, the values of the normal trace, $\langle F \cdot \nu|_{\partial\Omega}, \phi|_{\partial\Omega} \rangle$, depend only on the values of ϕ over $\partial\Omega$. In contrast, for $F \in \mathcal{DM}^{ext}(\Omega)$, the values of $\langle F \cdot \nu|_{\partial\Omega}, \phi|_{\partial\Omega} \rangle$ also depend, in principle, on the values of the first derivatives of ϕ over $\partial\Omega$, since $\phi|_{\partial\Omega}$ must be viewed as elements of $\text{Lip}(\gamma, \partial\Omega)$, for some $\gamma > 1$.

Finally, we establish the following useful product rule.

Theorem 3.2 (Product Rule). *Let $F = (F_1, \dots, F_N) \in \mathcal{DM}^*(D)$. Let $g \in C(D)$ be such that $\partial_{x_j} g(x)$ is $|F_j|$ -integrable, for each $j = 1, \dots, N$, and the set of non-Lebesgue points of $\partial_{x_j} g(x)$ has $|F_j|$ -measure zero. Then $gF \in \mathcal{DM}^*(D)$ and*

$$\text{div}(gF) = g \text{div} F + \nabla g \cdot F. \tag{3.13}$$

Proof. For any $\phi \in C_0^1(D)$, we have

$$\langle \text{div}(gF), \phi \rangle = -\langle F, g \nabla \phi \rangle = -\langle F, \nabla(g\phi) \rangle + \langle F, \phi \nabla g \rangle. \tag{3.14}$$

Therefore, it suffices to show that

$$\langle F, \nabla(g\phi) \rangle = -\langle \text{div} F, g\phi \rangle. \tag{3.15}$$

Set $g^\varepsilon = g * \omega^\varepsilon$. We have

$$\begin{aligned} -\langle \text{div} F, g^\varepsilon \phi \rangle &= \langle F, \nabla(g^\varepsilon \phi) \rangle = \sum_{j=1}^N \langle F_j, \partial_{x_j}(g^\varepsilon \phi) \rangle \\ &= \sum_{j=1}^N \{ \langle F_j, \phi \partial_{x_j} g^\varepsilon \rangle + \langle F_j, g^\varepsilon \partial_{x_j} \phi \rangle \}. \end{aligned} \tag{3.16}$$

Let $\varepsilon \rightarrow 0$. Then the right-hand side converges to

$$\sum_{j=1}^N \{ \langle F_j, \phi \partial_{x_j} g \rangle + \langle F_j, g \partial_{x_j} \phi \rangle \} = \langle F, \nabla(g\phi) \rangle,$$

by the assumption on the set of non-Lebesgue points of $\partial_{x_j} g$, while the left-hand side of (3.16) converges to $-\langle \operatorname{div} F, g\phi \rangle$ by the Dominated Convergence Theorem, which implies (3.15). Then (3.13) follows. \square

Remark 3.3. The continuity assumption of $g(x)$ in Theorem 3.2 can be relaxed. In particular, when $F \in \mathcal{DM}^\infty(D)$, it requires only that $g \in BV(D; \mathbb{R})$ to have $gF \in \mathcal{DM}^\infty(D)$.

Remark 3.4. The results in this section for the extended vector fields over \mathbb{R}^N extend to a general context over Riemannian manifolds. In particular, if $a_{ij}(x)$ are smooth functions on the open set $D \subset \mathbb{R}^N$ and the $N \times N$ matrix $(a_{ij})(x)$ is symmetric, positive definite, the results can easily be generalized to the extended vector fields $F = (F_1, \dots, F_N)$ in $L^p(D; \mathbb{R}^N)$, $1 \leq p \leq \infty$, or $M(D; \mathbb{R}^N)$, satisfying the condition:

$$\sum_{i,j} a_{ij} \partial_i F_j \in \mathcal{M}(D).$$

This is clear from the fact that no specific property of the Euclidean metric has been used in our analysis.

4. Applications to the Euler Equations for Gas Dynamics

In this section, as a direct application of the theory developed in Sect. 3, we establish the uniqueness and stability of Riemann solutions that may contain vacuum for the Euler equations for gas dynamics in Lagrangian coordinates.

Denote $\mathbb{R}_+^2 = (0, \infty) \times \mathbb{R}$ and $\overline{\mathbb{R}_+^2} = [0, \infty) \times \mathbb{R}$. We consider $\tau \in \mathcal{M}^+(\mathbb{R}_+^2)$ satisfying $\tau \geq c \mathcal{L}^2$ for some $c > 0$, where \mathcal{L}^k is the k -dimensional Lebesgue measure. Let $v \in L^\infty(\mathbb{R}_+^2)$ and $\tau_0 \in \mathcal{M}^+(\mathbb{R})$ with $\tau_0 \geq c \mathcal{L}^1$. We assume that τ , v , and τ_0 satisfy

$$\iint_{\mathbb{R}_+^2} (\phi_t \tau - v \phi_x dt dx) + \int_{\mathbb{R}} \phi(0, x) \tau_0(x) dx = 0, \tag{4.1}$$

for any $\phi \in C_0^1(\mathbb{R}^2)$.

Definition 4.1. Let τ and τ_0 be as above. We say that a function $\phi(t, x)$ defined on $\overline{\mathbb{R}_+^2}$ is a τ -test function if it satisfies the following:

1. $\operatorname{spt}(\phi)$ is a compact subset of $\overline{\mathbb{R}_+^2}$ and ϕ is continuous on \mathbb{R}_+^2 ;
2. ϕ_t and ϕ_x are τ -measurable; and ϕ_t is τ -integrable over \mathbb{R}_+^2 , that is, the integrals $\iint_{\mathbb{R}_+^2} (\phi_t)_{\pm} \tau$ exist and at least one of them is finite;
3. $\lim_{\substack{t \rightarrow 0 \\ x \rightarrow a}} \phi(t, x) = \phi(0, a)$ for τ_0 -a.e. $a \in \mathbb{R}$.

Theorem 4.1. *Let τ , v , and τ_0 be as above. Then*

1. *the nonnegative measure τ admits a slicing of the form $\tau = dt \otimes \mu_t(x)$ with $\mu_t \in \mathcal{M}^+(\mathbb{R})$ for \mathcal{L}^1 -a.e. $t > 0$. More precisely, for all $\phi \in C_0(\mathbb{R}_+^2)$,*

$$\iint \phi(t, x) \tau = \int \left(\int \phi(t, x) \mu_t(x) \right) dt;$$

2. *the points $(t, x) \in \mathbb{R}_+^2$ such that $\mu_t(x) > 0$, with the exception of a set of \mathcal{H}^1 -measure zero, form a countable union of vertical line segments, called vacuum lines. In particular, $\tau(l) = 0$ for any non-vertical straight line segment l ;*

3. *the identity (4.1) holds for any τ -test function $\phi(t, x)$.*

The proof of Theorem 4.1 is given in Sect. 5. As a corollary, we have

Corollary 4.1. *Let τ , v , and τ_0 be as above. Let $\bar{p}(t, x)$ be a nonnegative function over \mathbb{R}_+^2 , continuous on \mathbb{R}_+^2 , such that $\phi \bar{p}$ is a τ -test function for any $\phi \in C_0^1(\mathbb{R}^2)$, $\bar{p}_t \leq 0$, τ -a.e., and $\bar{p}_x \in L^1_{loc}(\mathbb{R}_+^2)$. Then, for any nonnegative function $\zeta \in C_0^1(\mathbb{R})$,*

$$\limsup_{t \rightarrow 0+} \int \zeta(x) \bar{p}(t, x) \mu_t(x) \leq \int \zeta(x) \bar{p}(0, x) \tau_0(x). \tag{4.2}$$

Proof. First we have from Theorem 4.1 that (4.1) holds for $\phi = \bar{p}\psi$ with $\psi \in C_0^1(\mathbb{R}^2)$. Then we choose $\psi = \psi^\varepsilon(t, x) := \sigma^\varepsilon(t - t_0)\zeta(x)$ with $\zeta(x) \geq 0$, $t_0 > 0$, and

$$\begin{aligned} 0 &\leq \sigma^\varepsilon \in C_0^1(\mathbb{R}), \\ \sigma^\varepsilon(t - t_0) &\rightarrow \chi_{(-\infty, t_0)}(t), \\ \delta^\varepsilon(t - t_0) &= -\frac{d}{dt} \sigma^\varepsilon(t - t_0) \rightarrow \delta_{t_0}, \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

where δ_{t_0} is the Dirac measure concentrated at t_0 , and the convergences are in $L^1(\mathbb{R})$ and $\mathcal{M}(\mathbb{R})$, respectively. We obtain from (4.1) that

$$\begin{aligned} 0 &= \iint_{t>0} ((\psi^\varepsilon \bar{p})_t \mu_t dt - (\psi^\varepsilon \bar{p})_x v dt dx) + \int (\psi^\varepsilon \bar{p}_0) \tau_0 \\ &= - \int \delta^\varepsilon \left(\int \zeta \bar{p} \mu_t \right) dt + \iint \bar{p}_t \psi^\varepsilon \mu_t dt - \iint (\psi^\varepsilon \bar{p})_x v dt dx + \int (\psi^\varepsilon \bar{p}_0) \tau_0. \end{aligned}$$

Now, using that $\bar{p}_t \leq 0$, τ -a.e., we obtain

$$- \int \delta^\varepsilon \left(\int \zeta \bar{p} \mu_t \right) dt + C_0 \iint_{\substack{0 \leq t \leq t_0 \\ x \in \text{spt}(\zeta)}} (|\bar{p}_x| + |\zeta'|) dt dx + \int (\psi^\varepsilon \bar{p}_0) \tau_0 \geq 0.$$

Assuming that t_0 is a Lebesgue point of $g(t) = \int \zeta \bar{p} \mu_t$ and letting $\varepsilon \rightarrow 0$ yield

$$\int \zeta \bar{p} \mu_{t_0} \leq C \iint_{\substack{0 \leq t \leq t_0 \\ x \in \text{spt}(\zeta)}} (|\bar{p}_x| + |\zeta'|) dt dx + \int \zeta \bar{p}_0 \tau_0. \tag{4.3}$$

Now, taking the lim sup as $t_0 \rightarrow 0$ in both sides of (4.3), we finally arrive at (4.2). \square

We now consider the solutions of the Euler equations (1.1)–(1.3) for gas dynamics in the sense of distributions such that τ is a nonnegative Radon measure, with $\tau \geq c\mathcal{L}^2$ for some $c > 0$, and $v(t, x)$ and $S(t, x)$ are bounded τ -measurable functions, along with our understanding that the constitutive relations (1.4) for $(\tau, p, e, \theta, S)(t, x)$ hold \mathcal{L}^2 -almost everywhere out of the vacuum lines, in the set where τ is absolutely continuous with respect to \mathcal{L}^2 , and both $p(t, x)$ and $e(t, x)$ are defined as zero on the remaining set with measure zero in \mathbb{R}_+^2 , including the vacuum lines.

We consider the Cauchy problem for (1.1)–(1.3):

$$(\tau, v, S)|_{t=0} = (\tau_0, v_0, S_0)(x), \tag{4.4}$$

where $\tau_0(x)$ is a nonnegative Radon measure over \mathbb{R} , $\tau_0 \geq c\mathcal{L}^1$ for some $c > 0$, $v_0(x)$ and $S_0(x)$ are bounded τ_0 -measurable functions, and $e_0(x) = e(\tau_0(x), S_0(x))$ a.e. out of the countable points $\{x_k\}$ such that $\tau_0(x_k) > 0$, the initial vacuum set.

Set $\Pi_T = (0, T) \times \mathbb{R}$ and $\Pi_T^* = (-\infty, T) \times \mathbb{R}$ for $T > 0$. Let D and F be functions or measures over Π_T . Let D_0 be a function or a measure over \mathbb{R} . By *weak formulation* on Π_T for the Cauchy problem:

$$D_t + F_x = 0, \tag{4.5}$$

$$D|_{t=0} = D_0, \tag{4.6}$$

we mean that, for a suitable set of test functions $\phi(t, x)$ defined on Π_T^* ,

$$\iint_{\Pi_T} (\phi_t D + \phi_x F) + \int_{\mathbb{R}} \phi(0, x) D_0 = 0. \tag{4.7}$$

Analogously, if the identity “=” in (4.5) is replaced by “ \geq ” or “ \leq ”, the weak formulation of the corresponding problem (4.5) and (4.6) is (4.7) with “=” replaced by “ \leq ” or “ \geq ”, respectively, for a suitable set of *nonnegative* test functions defined on Π_T^* .

Denote $W = (\tau, v, S)$, $f(W) = (-v, p(\tau, S), 0)$, $\eta(W) = e(\tau, S) + \frac{v^2}{2}$, $q(W) = vp(\tau, S)$, and

$$\begin{aligned} \alpha(W, \bar{W}) &= \eta(W) - \eta(\bar{W}) - \nabla\eta(\bar{W}) \cdot (W - \bar{W}), \\ \beta(W, \bar{W}) &= q(W) - q(\bar{W}) - \nabla\eta(\bar{W}) \cdot (f(W) - f(\bar{W})). \end{aligned}$$

Observe that $\nabla\eta(\bar{W}) = (-\bar{p}, -\bar{v}, \bar{\theta})$.

Definition 4.2. *We say that $W(t, x)$ is a distributional entropy solution of (1.1)–(1.3), and (4.4) in Π_T if τ is a Radon measure on Π_T with $\tau \geq c\mathcal{L}^2$ for some $c > 0$, v and S are bounded τ -measurable functions such that the weak formulation of (1.1)–(1.3), (1.7), and (4.4) is satisfied for all test functions in $C_0^1(\Pi_T^*)$, and $S(t, \cdot) \rightharpoonup S_0(\cdot)$, as $t \rightarrow 0$, in the weak-star topology of $L^\infty(\mathbb{R})$.*

Observe that the weak formulation implies that $\mu_t \rightharpoonup \tau_0$ in $\mathcal{M}(\mathbb{R})$, and $v(t, \cdot) \rightharpoonup v_0(\cdot)$ and $E(t, \cdot) \rightharpoonup E_0(\cdot)$ in the weak-star topology of $L^\infty(\mathbb{R})$, as $t \rightarrow 0$, where $E = e + v^2/2$. We also remark that these convergences can be strengthened to the convergences in $L^1_{loc}(\mathbb{R})$ in the case that τ is a bounded measurable function, as an easy consequence of the \mathcal{DM}^∞ theory (cf. [5]).

As shown by Wagner [25], by means of the transformation from Eulerian to Lagrangian coordinates, bounded measurable entropy solutions of the Euler equations

in Eulerian coordinates transform into distributional entropy solutions of (1.1)–(1.3), (1.7) and (4.4), satisfying the additional restriction that the weak formulation of (1.2), (1.3), and (1.7) holds for test functions with compact support in Π_T such that $\phi_t = g$, $\phi_x = h \tau$, where $g, h \in L^\infty(\Pi_T, \tau)$. It is also shown through an example in [25] that distributional entropy solutions without the additional restriction may have no physical meaning.

Now we consider the Riemann solution $\bar{W}(t, x)$ associated with the Riemann problem for (1.1)–(1.3) with initial condition:

$$\bar{W}_0(x) = \begin{cases} W_L, & x < 0, \\ W_R, & x > 0, \end{cases} \tag{4.8}$$

where W_L and W_R are two constant states in the physical domain $\{W = (\tau, v, S) : \tau > 0\}$. First, we address the case that $\bar{W}(t, x)$ is a bounded self-similar entropy solution of (1.1)–(1.3) which consists of at most two rarefaction waves, one corresponding to the first characteristic family and another corresponding to the third one, and possibly one contact discontinuity on the line $x = 0$. Then, $\bar{W}(t, x)$ has the following general form:

$$\bar{W}(x, t) = \begin{cases} W_L, & x/t < \xi_1, \\ R_1(x/t), & \xi_1 \leq x/t < \xi_2, \\ W_M, & \xi_2 \leq x/t < 0, \\ W_N, & 0 < x/t < \xi_3, \\ R_3(x/t), & \xi_3 \leq x/t < \xi_4, \\ W_R, & x/t \geq \xi_4. \end{cases} \tag{4.9}$$

In what follows we use the notation $W^p = (\tau, S)$ and $\bar{W}^p = (\bar{\tau}, \bar{S})$.

Theorem 4.2. *Let $\bar{W}(t, x)$ be the Riemann solution (4.9), and let $W(t, x)$ be any distributional entropy solution of (1.1)–(1.3) and (4.4) with $W_0 \in L^\infty(\mathbb{R}; \mathbb{R}^3)$. Then there exist positive constants C and K_0 , and a function $\omega \in L^\infty(\Pi_T)$, positive a.e. in Π_T , such that, for any $X > 0$ and a.e. $t > 0$,*

$$\begin{aligned} & \int_{|x| \leq X} \left(|v(t, x) - \bar{v}(t, x)|^2 + |W_{a.c.}^p(t, x) - \bar{W}^p(t, x)|^2 \omega(t, x) \right) dx \\ & \leq C \int_{|x| \leq X + K_0 t} \left(|v_0(x) - \bar{v}_0(x)|^2 + |W_0^p(x) - \bar{W}_0^p(x)|^2 \omega(0, x) \right) dx. \end{aligned} \tag{4.10}$$

Proof. We divide the proof into four steps.

Step 1. Consider the measure

$$\varpi := \alpha(W, \bar{W})_t + \beta(W, \bar{W})_x.$$

Given any $X > 0$ and $t > 0$, let $t_0 \in (0, t)$ and

$$\Omega_{t_0, t} = \{(\sigma, x) : |x| < X + K_0(t - \sigma), t_0 < \sigma < t\},$$

for $K_0 > 0$ to be suitably chosen later.

First, by the Gauss-Green formula (Theorem 3.1), we have

$$\varpi(\Omega_{t_0,t}) = \langle (\alpha, \beta) \cdot \nu |_{\partial\Omega_{t_0,t}}, 1 \rangle.$$

Now, let $\zeta_i, i = 1, \dots, 4$, be nonnegative functions in $C_0^\infty(\mathbb{R}^2)$ such that

$$\begin{aligned} \sum_{i=1}^4 \zeta_i &= 1, & \text{on } \partial\Omega_{t_0,t}, \\ \zeta_1 &= 1, & \text{on } \{(0, x) : |x| < X + K_0(t - t_0)\}, \\ \zeta_2 &= 1, & \text{on } \{(t, x) : |x| < X\}, \\ (\text{spt}(\zeta_3) \cup \text{spt}(\zeta_4)) \cap ((\{t\} \times \mathbb{R}) \cup (\{t_0\} \times \mathbb{R})) &= \emptyset. \end{aligned}$$

We choose ζ_3 and ζ_4 so that $\text{spt}(\zeta_3)$ intersects the left lateral side of $\Omega_{t_0,t}$ but not the right, and $\text{spt}(\zeta_4)$ intersects its right lateral side but not the left. We have

$$\varpi(\Omega_{t_0,t}) = \langle (\alpha, \beta) \cdot \nu |_{\partial\Omega_{t_0,t}}, \zeta_1 + \zeta_2 + \zeta_3 + \zeta_4 \rangle.$$

In what follows, we will use the notations:

$$\hat{g} = \int_0^1 (1 - \sigma)g(\sigma\tau + (1 - \sigma)\bar{\tau}, \sigma S + (1 - \sigma)\bar{S}) d\sigma$$

for any function $g = g(\tau, S)$, and

$$\tilde{g} = \int_0^1 g(\sigma\tau + (1 - \sigma)\bar{\tau}, \sigma S + (1 - \sigma)\bar{S}) d\sigma.$$

Step 2. We now prove the following five estimates.

1. $\langle (\alpha, \beta) \cdot \nu |_{\partial\Omega_{t_0,t}}, \zeta_3 \rangle \geq 0$ and $\langle (\alpha, \beta) \cdot \nu |_{\partial\Omega_{t_0,t}}, \zeta_4 \rangle \geq 0$.

Indeed, let $z_* = (t_*, x_*)$ be the center of $\Omega_{t_0,t}$. We consider the following deformation of $\partial\Omega_{t_0,t}$:

$$\Psi(z, s) := z + \varepsilon s(z_* - z), \quad z \in \partial\Omega_{t_0,t}, \quad s \in [0, 1],$$

where ε is chosen so small that

$$\begin{aligned} &(\text{spt}(\zeta_3) \cup \text{spt}(\zeta_4)) \\ &\cap (\{(t - \varepsilon s(t - t_*), x) : x \in \mathbb{R}\} \cup \{(t_0 + \varepsilon s(t_* - t_0), x) : x \in \mathbb{R}\}) = \emptyset, \end{aligned}$$

for $s \in [0, 1]$. Then

$$\langle (\alpha, \beta) \cdot \nu |_{\partial\Omega_{t_0,t}}, \zeta_3 \rangle = \lim_{s \rightarrow 0} \frac{1}{s} \iint_{\Psi(\partial\Omega_{t_0,t} \times (0,s))} (K_0\alpha - \beta) |\nabla h| \zeta_3 d\sigma dx \geq 0,$$

by choosing K_0 such that $K_0\alpha \geq \beta$, which is possible by what follows.

2. Given $\delta > 0$ and bounded sets $B_1 \subset V_\delta := \{(\tau, v, S) : \tau > \delta\}$ and $B_2 \subset \mathbb{R}^2$, there exists a constant $K_0 = K_0(\delta, B_1, B_2) > 0$ such that $K_0\alpha(W, \bar{W}) \geq \beta(W, \bar{W})$, for any $\bar{W} \in B_1$ and $W \in V_\delta$ with $(v, S) \in B_2$.

In fact, we first have

$$\begin{aligned} \alpha(W, \bar{W}) &= \frac{1}{2}(v - \bar{v})^2 + \hat{e}_{\tau\tau}(\tau - \bar{\tau})^2 + 2\hat{e}_{\tau S}(\tau - \bar{\tau})(S - \bar{S}) + \hat{e}_{SS}(S - \bar{S})^2, \\ \beta(W, \bar{W}) &= (v - \bar{v})(p - \bar{p}). \end{aligned}$$

If $W(t, x)$ and $\overline{W}(t, x)$ belong to a bounded set B in V_δ , we can find K_0 depending only on B such that $K_0\alpha \geq \beta$. Now, since $(\bar{\tau}, \bar{v}, \bar{S}) \in B_1$ and $(v, S) \in B_2$, it suffices to show that, for τ sufficiently large, we have $\beta(W, \overline{W}) \leq \alpha(W, \overline{W})$. Notice that

$$\begin{aligned} \beta &\leq \frac{1}{2}(v - \bar{v})^2 + \frac{1}{2}\tilde{p}_\tau^2(\tau - \bar{\tau})^2 + \tilde{p}_\tau \tilde{p}_S(\tau - \bar{\tau})(S - \bar{S}) + \frac{1}{2}\tilde{p}_S^2(S - \bar{S})^2 \\ &\leq \frac{1}{2}(v - \bar{v})^2 + \tilde{p}_\tau^2(\tau - \bar{\tau})^2 + \tilde{p}^2(S - \bar{S})^2. \end{aligned}$$

On the other hand,

$$\alpha \geq \frac{1}{2}(v - \bar{v})^2 + \frac{\hat{e}_{\tau\tau}}{2(K+1)}(\tau - \bar{\tau})^2 + \frac{1}{2} \left(\hat{e}_{SS} - \frac{K+1}{K} \widehat{\left(\frac{e_{\tau S}^2}{e_{\tau\tau}} \right)} \right) (S - \bar{S})^2,$$

for any $K > 0$. Now, \tilde{p}_τ^2 decays faster than $\hat{e}_{\tau\tau} = -\hat{p}_\tau$ as $\tau \rightarrow \infty$, \tilde{p}_S^2 decays faster than \hat{e}_{SS} as $\tau \rightarrow \infty$, and, for K sufficiently large,

$$\hat{e}_{SS} - \frac{K+1}{K} \widehat{\left(\frac{e_{\tau S}^2}{e_{\tau\tau}} \right)} > c \hat{e}_{SS},$$

for some $c > 0$ sufficiently small, since $\gamma > 1$, as one can easily check from (1.4).

Hence, $\beta(W, \overline{W}) \leq \alpha(W, \overline{W})$ for τ larger than a certain τ_* . Then the assertion follows.

3. Similarly, we have $\langle (\alpha, \beta) \cdot \nu |_{\partial\Omega_{t_0,t}}, \zeta_4 \rangle \geq 0$.

4. As for ζ_2 , we have

$$\begin{aligned} &\langle (\alpha, \beta) \cdot \nu |_{\partial\Omega_{t_0,t}}, \zeta_2 \rangle \\ &= \lim_{s \rightarrow 0} \frac{1}{s} \iint_{\Psi(\partial\Omega_{t_0,t} \times (0,s))} (\alpha, \beta) \cdot \nabla h \zeta_2 d\sigma dx \\ &\geq \lim_{s \rightarrow 0} \frac{1}{\varepsilon(t-t_*)s} \int_{t-\varepsilon s(t-t_*)}^t d\sigma \int_{|x| \leq X-\varepsilon K_1 \frac{T}{2}} \{ \eta(W) - \eta(\overline{W}) - \bar{v}(v - \bar{v}) \\ &\quad + \bar{p}(\tau_{\text{a.c.}} - \bar{\tau}) + \bar{\theta}(S - \bar{S}) \} dx \\ &= \int_{\substack{|x| \leq X-\varepsilon K_1 \frac{T}{2} \\ \sigma=t}} \{ \eta(W_{\text{a.c.}}) - \eta(\overline{W}) - \bar{v}(v - \bar{v}) + \bar{p}(\tau_{\text{a.c.}} - \bar{\tau}) + \bar{\theta}(S - \bar{S}) \} dx \\ &= \int_{\substack{|x| \leq X-\varepsilon K_1 \frac{T}{2} \\ \sigma=t}} \left(\frac{1}{2}|v - \bar{v}|^2 + Q(W_{\text{a.c.}}^p(t, x) - \overline{W}^p(t, x)) \right) dx, \end{aligned}$$

if t is a Lebesgue point of

$$g(s) = \int_{|x| \leq X-\varepsilon K_1 \frac{T}{2}} \alpha(W_{\text{a.c.}, \overline{W}})(s, x) dx,$$

where Q is the quadratic form associated with the symmetric matrixe

$$A = \begin{bmatrix} \hat{e}_{\tau\tau} & \hat{e}_{\tau S} \\ \hat{e}_{\tau S} & \hat{e}_{SS} \end{bmatrix}.$$

5. For ζ_1 , we have

$$\begin{aligned} & \langle (\alpha, \beta) \cdot \nu |_{\partial\Omega_{t_0,t}}, \zeta_1 \rangle \\ &= \lim_{s \rightarrow 0} \frac{1}{s} \iint_{\Psi(\partial\Omega_{t_0,t} \times (0,s))} (\alpha, \beta) \cdot \nu \zeta_1 |\nabla h| d\sigma dx \\ &\geq - \lim_{s \rightarrow 0} \frac{1}{s} \int_{t_0}^{t_0 + \varepsilon s(t_* - t_0)} \int_{|x| \leq X + K_0(t-t_0)} \alpha(W, \bar{W}) \frac{1}{\varepsilon(t_* - t_0)} d\sigma dx \\ &= - \int_{\substack{|x| \leq X + K_0(t-t_0) \\ \sigma = t_0}} \{ \eta(W) - \eta(\bar{W}) - \bar{v}(v - \bar{v}) + \bar{p}(\mu_{t_0} - \bar{\tau}) - \bar{\theta}(S - \bar{S}) \}, \end{aligned}$$

where we have also used that $\mu_\sigma \rightarrow \mu_{t_0}$ as $\sigma \rightarrow t_0 + 0$, for a.e. $t_0 > 0$, with μ_t as in Theorem 4.1, and that \bar{p} is continuous on $[t_0, t] \times \mathbb{R}$.

Step 3. On the other hand,

$$\begin{aligned} \varpi(\Omega_{t_0,t}) &= \sum_{i=1}^4 \varpi(\tilde{l}_i \cap \Omega_{t_0,t}) + \varpi(l \cap \Omega_{t_0,t}) + \varpi(\Omega_1 \cap \Omega_{t_0,t}) \\ &\quad + \varpi(\Omega_3 \cap \Omega_{t_0,t}) + \varpi(\Omega_{t_0,t} - (\cup_{i=1}^4 \tilde{l}_i \cup l \cup \Omega_1 \cup \Omega_3)), \end{aligned}$$

where Ω_1 and Ω_3 are the left and right rarefaction regions, $\tilde{l}_i, 1 \leq i \leq 4$, are the lines bounding the rarefaction regions Ω_1 and Ω_3 , and l is the line $\{x = 0\}$, where $\bar{W}(t, x)$ has a contact discontinuity.

We first observe that, on $\Omega_{t_0,t} - (\cup_{i=1}^4 \tilde{l}_i \cup l \cup \Omega_1 \cup \Omega_3)$, the measure ϖ reduces to $-\bar{\theta} \partial_t S$ which is nonpositive. Now, we have

$$\varpi = -\text{div}(F_1 + F_2 + F_3),$$

where

$$F_1 = \bar{v}(v - \bar{v}, p - \bar{p}), \quad F_2 = -\bar{p}(\tau - \bar{\tau}, \bar{v} - v), \quad F_3 = \bar{\theta}(S - \bar{S}, 0),$$

and $\text{div} := \text{div}_{t,x}$. Applying the product rule (Theorem 3.2), we get

$$\text{div} F_1 = \bar{v}_t(v - \bar{v}) + \bar{v}_x(p - \bar{p}), \quad \text{div} F_2 = -\bar{p}_t(\tau - \bar{\tau}) + \bar{p}_x(v - \bar{v}).$$

Hence,

$$\text{div} F_1(\tilde{l}_j \cap \Omega_{t_0,t}) = \text{div} F_1(l \cap \Omega_{t_0,t}) = 0, \quad j = 1, \dots, 4,$$

since $\text{div} F_1$ is absolutely continuous with respect to \mathcal{L}^2 . Also,

$$\text{div} F_3(\tilde{l}_j \cap \Omega_{t_0,t}) \geq 0, \quad j = 1, \dots, 4.$$

On the other hand, since $F_3 \in \mathcal{DM}^\infty(\Omega_{t_0,t})$ and $\nu|_l = (0, 1)$, we have

$$\operatorname{div} F_3(l \cap \Omega_{t_0,t}) = [\langle F_3 \cdot \nu|_l, 1 \rangle] = 0,$$

where the bracket denotes the difference between the normal traces from the right and the left, which make sense for $F_3 \in \mathcal{DM}^\infty$ because the normal traces of \mathcal{DM}^∞ fields are functions in L^∞ over the boundaries.

Concerning F_2 , we have

$$\operatorname{div} F_2(\tilde{l}_j \cap \Omega_{t_0,t}) = 0, \quad j = 1, \dots, 4,$$

since \bar{p}_t is τ -integrable and $\tau(\tilde{l}_j) = 0$, $j = 1, \dots, 4$. On the other hand, \bar{p}_t vanishes on l so that

$$\operatorname{div} F_2(l \cap \Omega_{t,t_0}) = 0.$$

Finally, using the product rule (Theorem 3.2), the fact that $W(t, x)$ and $\bar{W}(t, x)$ are distributional solutions of (1.1)–(1.3) and (4.4), and $\bar{S}(t, x)_t = 0$, we obtain, for $j = 1, 3$,

$$\begin{aligned} & \varpi(\Omega_j \cap \Omega_{t_0,t}) \\ &= - \iint_{\Omega_j \cap \Omega_{t_0,t}} \{(\bar{v}(v - \bar{v}))_t + (\bar{v}(p - \bar{p}))_x - (\bar{p}(\tau - \bar{\tau}))_t + (\bar{p}(v - \bar{v}))_x + (\bar{\theta}(S - \bar{S}))_t\} \\ &= - \iint_{\Omega_j \cap \Omega_{t_0,t}} \{\bar{v}_t(v - \bar{v}) + \bar{v}_x(p - \bar{p}) - \bar{p}_t(\tau - \bar{\tau}) + \bar{p}_x(v - \bar{v}) + \bar{\theta}_t(S - \bar{S}) + \bar{\theta}_t S_t\} \\ &\leq - \iint_{\Omega_j \cap \Omega_{t_0,t}} \{\bar{v}_x(p - \bar{p}) - \bar{p}_\tau \bar{v}_x(\tau - \bar{\tau}) + \bar{\theta}_\tau \bar{v}_x(S - \bar{S})\} \\ &\leq - \iint_{\Omega_j \cap \Omega_{t_0,t}} \bar{v}_x(p - \bar{p} - \bar{p}_\tau(\tau_{a.c.} - \bar{\tau}) - \bar{p}_S(S - \bar{S})) dx ds + \iint_{\Omega_j \cap \Omega_{t_0,t}} \bar{v}_x \bar{p}_\tau \tau_{sing} \leq 0, \end{aligned} \tag{4.11}$$

since $\bar{p}_\tau < 0$, \bar{v}_x is bounded, $\bar{v}_x(t, x) \geq 0$ everywhere over Ω_1 and Ω_3 , and $\nabla_{W,p}^2 p > 0$.

Step 4. Putting all these estimates together, we have

$$\begin{aligned} & \int_{|x| \leq X} \left(|v(t, x) - \bar{v}(t, x)|^2 + |W_{a.c.}(t, x) - \bar{W}(t, x)|^2 \omega(t, x) \right) dx \\ & \leq 2 \int_{\substack{|x| \leq X + K_0(t-t_0) \\ \sigma=t_0}} \{ \eta(W) - \eta(\bar{W}) - \bar{v}(v - \bar{v}) + \bar{p}(\mu_{t_0} - \bar{\tau}) - \bar{\theta}(S - \bar{S}) \}. \end{aligned}$$

Now, applying Corollary 4.1, we finally get (4.10). \square

Corollary 4.2. *Let $W(t, x)$ and $\bar{W}(t, x)$ satisfy the conditions of Theorem 4.2 and $W_0(x) = \bar{W}_0(x)$. Then $\tau(t, x)$ is absolutely continuous with respect to \mathcal{L}^2 in Π_T and $W(t, x) = \bar{W}(t, x)$ a.e. in Π_T .*

Proof. From Theorem 4.2, we have

$$W_{a.c.}(t, x) = \overline{W}(t, x), \quad \text{a.e. in } \Pi_T.$$

Hence, τ_{sing} must satisfy the weak formulation of

$$\tau_t = 0, \quad \tau|_{t=0} = 0,$$

for the test functions in $C_0^1(\Pi_T^*)$. In particular, there exists $y \in BV_{loc}(\Pi_T)$ such that

$$\partial_x y = \tau_{sing}, \quad \partial_t y = 0.$$

Therefore,

$$\tau_{sing} = dt \otimes v_t(x),$$

where $v_t(\cdot) = \frac{dy}{dx}(t, \cdot)$, a.e. $t \in (0, T)$. Since y is independent of t , we have that v_t is also independent of t , say, $v_t(x) = v_0(x)$ and

$$\tau_{sing} = dt \otimes v_0.$$

Furthermore, since $v_t \rightarrow 0$ as $t \rightarrow 0$, we conclude

$$v_0 \equiv 0.$$

□

We now consider the case that the Riemann solution, with the initial condition (4.8), has a vacuum line at $x = 0$. In this case, the Riemann solution $\overline{W}(t, x)$ has the following form:

$$\overline{W}(x, t) = \begin{cases} W_L, & x/t < \xi_L, \\ R_1(x/t), & \xi_L \leq x/t < 0, \\ ((\bar{v}_1 + \bar{v}_2)/2, (\bar{v}_2 - \bar{v}_1)t dt \otimes \delta_0(x), (\bar{S}_L + \bar{S}_R)/2), & x = 0, \\ R_3(x/t), & 0 < x/t \leq \xi_R, \\ W_R, & x/t > \xi_R. \end{cases} \tag{4.12}$$

Here $R_1(x/t)$ and $R_3(x/t)$ are as above the rarefaction waves of the first and third characteristic families, respectively, $\bar{v}_1 = \lim_{\xi \rightarrow 0^-} \bar{v}(\xi)$, $\bar{v}_2 = \lim_{\xi \rightarrow 0^+} \bar{v}(\xi)$, and $\delta_0(x)$ is the Dirac measure over \mathbb{R} concentrated at $x = 0$. It is easy to check that $\overline{W}(t, x)$ is a distributional solution of (1.1)–(1.3) and (4.4). The values of \bar{v} and \bar{S} on the line $x = 0$ could be taken as any other constants instead of $\frac{\bar{v}_1 + \bar{v}_2}{2}$ and $\frac{\bar{S}_L + \bar{S}_R}{2}$, respectively, while the formula of $\bar{\tau}$ at $x = 0$, $(\bar{v}_2 - \bar{v}_1)t dt \otimes \delta_0(x)$, is dictated by the fact that (1.1) must hold in the sense of distributions.

Theorem 4.3. *Let $\overline{W}(t, x)$ be a Riemann solution containing vacuum as described in (4.12). Let $W(t, x)$ be a distributional entropy solution of (1.1)–(1.3) and (4.4) in Π_T with $W_0 \in L^\infty(\mathbb{R}; \mathbb{R}^3)$. Then there exist positive constants C and K_0 , and a function $\omega \in L^\infty(\Pi_T)$, positive a.e. in Π_T , such that, for all $X > 0$ and a.e. $t > 0$,*

$$\begin{aligned} & \int_{|x| \leq X} \left(|v(t, x) - \bar{v}(t, x)|^2 + |W_{a.c.}^p(t, x) - \overline{W}^p(t, x)|^2 \omega(t, x) \right) dx \\ & \leq C \int_{|x| \leq X + K_0 t} \left(|v_0(x) - \bar{v}_0(x)|^2 + |W_0^p(x) - \overline{W}_0^p(x)|^2 \omega(0, x) \right) dx. \end{aligned} \tag{4.13}$$

Proof. Let F_1 , F_2 , and F_3 be as in the proof of Theorem 4.2. We observe that, since $\bar{p} = 0$ for $x = 0$, we have

$$F_2 = -\bar{p}(\tau - \bar{\tau}_{a.c.}, \bar{v} - v),$$

so that the analysis for F_2 remains the same. Also, nothing needs to be changed concerning F_3 . As for $F_1 = \bar{v}(v - \bar{v}, p - \bar{p})$, we have a new aspect which is the fact that $\bar{v}(t, x)$ is discontinuous at l . Then we have

$$\operatorname{div} F_1(l_{t_0,t}) = [\langle F_1 \cdot \nu |_{l_{t_0,t}}, 1 \rangle],$$

where $l_{t_0,t} = l \cap \Omega_{t_0,t}$ and again $\nu|_l = (0, 1)$. Let $p_-(t, x)$ and $p_+(t, x)$ denote the functions in $L^\infty(l)$, given by the theory of \mathcal{DM}^∞ fields developed in [5], such that

$$\begin{aligned} \langle p_-, \zeta \rangle &= \langle (v, p) \cdot \nu|_l, \zeta \rangle_l, \\ \langle p_+, \zeta \rangle &= \langle (v, p) \cdot (-\nu)|_l, \zeta \rangle_l, \end{aligned} \quad \text{for any } \zeta \in C_0(l).$$

Hence, we have

$$\operatorname{div} F_1(l_{t_0,t}) = v_2 \int_{t_0}^t p_+(s) ds - v_1 \int_{t_0}^t p_-(s) ds.$$

On the other hand,

$$p_+(s) = p_-(s), \quad a.e. s > 0, \quad \text{and} \quad v_2 > v_1,$$

where the first follows from (1.2), and the second is a consequence of the construction of the Riemann solution containing vacuum. Therefore, we conclude

$$\operatorname{div} F_1(l_{t_0,t}) \geq 0.$$

The remainder of the proof follows exactly as in Theorem 4.2. \square

Again, we have the following corollary.

Corollary 4.3. *Let $\bar{W}(t, x)$ and $W(t, x)$ satisfy the hypotheses of Theorem 4.3 and $W_0(x) = \bar{W}_0(x)$. Then $(v, S)(t, x) = (\bar{v}, \bar{S})(t, x)$, \mathcal{L}^2 -a.e. in Π_T , and $\tau = \bar{\tau}$ in $\mathcal{M}(\Pi_T)$.*

Proof. From Theorem 4.3, we deduce that $W_{a.c}(t, x) = \bar{W}_{a.c}(t, x)$, \mathcal{L}^2 -a.e. in Π_T . Thus, as in the proof of Corollary 4.2, we deduce that τ_{sing} must be concentrated at $\{x = 0\}$. Then τ_{sing} must be equal to $(\bar{v}_2 - \bar{v}_1)t dt \otimes \delta_0(x)$ as a consequence of (1.1) in the sense of distributions. \square

5. Proof of Theorem 4.1

In this section, we give a detailed proof of Theorem 4.1. The arguments are strongly motivated by those in Wagner [25].

Proof. We divide the proof into ten steps.

Step 1. There exists $y \in BV_{loc}([0, \infty) \times \mathbb{R})$ such that

$$\partial_x y = \tau, \quad \partial_t y = v, \tag{5.1}$$

in the sense of distributions for $t > 0$.

Indeed, let ω^ε be a positive symmetric mollifier in \mathbb{R}^2 , and set $\tau^\varepsilon = \tau * \omega^\varepsilon$ and $v^\varepsilon = v * \omega^\varepsilon$, where we have extended τ and v as zero for $t < 0$. Define

$$y^\varepsilon(t, x) = \int_0^x \tau^\varepsilon(t, s) ds + \int_0^t v^\varepsilon(\sigma, 0) d\sigma. \tag{5.2}$$

Then $y^\varepsilon \in BV \cap C^1(\mathbb{R}_+^2)$. We easily check that $y^\varepsilon(t, x)$ satisfies

$$\partial_x y^\varepsilon = \tau^\varepsilon, \quad \partial_t y^\varepsilon = v^\varepsilon, \quad t > \varepsilon.$$

Now, $\|y^\varepsilon\|_{BV(\Omega)} \leq M_\Omega$, for any open set $\Omega \subset \subset \mathbb{R}_+^2$, where M_Ω is a positive constant independent of ε . Hence, by the compact embedding of $BV(\Omega)$ into $L^1(\Omega)$, there exists $y \in BV_{loc}(\mathbb{R}_+^2)$ such that

$$y^\varepsilon(t, x) \rightarrow y(t, x), \quad \text{in } L^1_{loc}(\mathbb{R}_+^2),$$

by passing to a subsequence if necessary. Clearly, $y(t, x)$ satisfies (5.1) in the sense of distributions in \mathbb{R}_+^2 .

Step 2. The measure $\tau(t, x)$ admits a slicing of the form $\tau = dt \otimes \mu_t$, where, for \mathcal{L}^1 -a.e. $t > 0$, $\mu_t \in \mathcal{M}^+(\mathbb{R})$.

Let $y(t, x)$ be a solution of (5.1). Since $y \in BV_{loc}(\mathbb{R}_+^2)$, then, for a.e. $t > 0$, $y(t, \cdot) \in BV_{loc}(\mathbb{R})$. Hence,

$$\tau = dt \otimes \frac{dy}{dx}(t, \cdot).$$

Step 3. The points $(t, x) \in \mathbb{R}_+^2$ such that $\mu_t(x) > 0$, with the possible exception of a set of \mathcal{H}^1 -measure zero, form a countable union of vertical line segments.

Again, since $y \in BV_{loc}(\mathbb{R}_+^2)$, then, for a.e. $x \in \mathbb{R}$, $y(\cdot, x) \in BV_{loc}(\mathbb{R}_+)$. Hence, $\partial_t y$ admits a slicing as

$$\partial_t y = \frac{dy}{dt}(\cdot, x) \otimes dx = v(\cdot, x) dt \otimes dx.$$

That is, for a.e. $x \in \mathbb{R}$, $y(\cdot, x)$ is a Lipschitz function on $[0, \infty)$ whose derivative is $v(\cdot, x)$. On the other hand, the jump set of $y(t, x)$, with the possible exception of a set of \mathcal{H}^1 -measure zero, is a countable union of C^1 curves $\{l_k\}_{k \in \mathbb{N}}$, by the structure theory of BV functions (see, e.g., [12, 13]). We then conclude that the lines l_k must be vertical, because, otherwise, we would have a subset $A \subset \mathbb{R}$ of positive measure such that, for $x \in A$, $\frac{dy}{dt}(\cdot, x)$ would be a singular measure, rather than an L^∞ function $v(\cdot, x)$, which proves the assertion.

Observe that $\mu_t \rightharpoonup \tau_0$ as $t \rightarrow 0+$, which follows from standard arguments by choosing suitable test functions in (4.1).

Step 4. Let $x_0 \in \mathbb{R}$ be such that $\tau_0(x_0) = \mu_t(x_0) = 0$, a.e. $t > 0$, and x_0 is a Lebesgue point of $\int_0^t v(\sigma, \cdot) d\sigma$, for all rational $t > 0$ and hence all fixed $t > 0$. Then there exists a solution of (5.1) satisfying

$$\lim_{t \rightarrow 0^+} y(t, x) = \int_{x_0}^x \tau_0, \quad \text{for a.e. } x \in \mathbb{R}. \tag{5.3}$$

Consider a mollifier ω^ε , with $\varepsilon = (\varepsilon_1, \varepsilon_2)$, of the form $\omega^\varepsilon(t, x) = \delta^{\varepsilon_1}(t)\delta^{\varepsilon_2}(x)$, where δ^{ε_j} , $j = 1, 2$, are standard positive symmetric mollifiers in \mathbb{R} . If $y^\varepsilon(t, x)$ is defined by (5.2), with 0 replaced by x_0 , then sending ε_1 to 0 first and next $\varepsilon_2 \rightarrow 0$ yield

$$\lim_{\varepsilon_2 \rightarrow 0} \lim_{\varepsilon_1 \rightarrow 0} y^\varepsilon(t, x) = \int_{x_0}^x \mu_t + \int_0^t v(\sigma, x_0) d\sigma, \quad \text{a.e. } (t, x) \in \mathbb{R}_+^2.$$

Hence,

$$y(t, x) = \int_{x_0}^x \mu_t + \int_0^t v(\sigma, x_0) d\sigma$$

provides the desired solution because of Step 4.

Now, since, for each $t \geq 0$, $y(t, x)$ is a strictly increasing function of x , there exists a well-defined monotone increasing continuous function $x(t, y)$ such that

$$x(t, y(t, x)) = x, \quad (t, x) \in \mathbb{R}_+^2, \tag{5.4}$$

where we have defined $y(t, x)$ on the vacuum lines by, say, $(y(t, x + 0) + y(t, x - 0))/2$.

Let Q and T be the transformations of \mathbb{R}_+^2 , $Q(t, x) = (t, y(t, x))$ and $T(t, y) = (t, x(t, y))$ so that $T(Q(t, x)) = (t, x)$. Observe that, given an open rectangle $R = (t_1, t_2) \times (a, b) \subset \mathbb{R}_+^2$, if a and b are such that $y(t, a)$ and $y(t, b)$ are continuous functions of t , then $Q(R)$ is an open set in \mathbb{R}_+^2 , which implies that T is continuous in \mathbb{R}_+^2 . Let

$$T_0(x) = \lim_{t \rightarrow 0^+} y(t, x), \quad Q_0(y) = \lim_{t \rightarrow 0^+} x(t, y).$$

Step 5. $T_{\#}\mathcal{L}^2 = \tau$ and $T_{0\#}\mathcal{L}^1 = \tau_0$, with notation from [13].

In fact, for a.e. $t > 0$, $y(t, x)$ is a strictly increasing BV_{loc} function of the variable x , and $\tau = dt \otimes \mu_t(x)$ with $\mu_t(x) = \frac{dy}{dx}(t, \cdot)$. Now, $T(t, y) = (t, x(t, y))$ and, for any open interval (a, b) and for each fixed $t > 0$,

$$x(t, \cdot)^{-1}(a, b) = (y(t, a + 0), y(t, b - 0)) = \tau(a, b).$$

Hence, the measure $T_{\#}\mathcal{L}^2$ equals $dt \otimes \tilde{\mu}_t$, where $\tilde{\mu}_t$ is the Stieltjes measure associated with $y(t, x)$, that is, $\tilde{\mu}_t = \mu_t$, which implies $T_{\#}\mathcal{L}^2 = \tau$. Similarly, we obtain $T_{0\#}\mathcal{L}^1 = \tau_0$.

Step 6. The map T is proper and onto.

The fact that T is onto follows from the fact that it is the inverse of Q which is defined everywhere in \mathbb{R}_+^2 . Now, since $Q(t, x)$ is a continuous function of t , for a.e. $x \in \mathbb{R}$, given any compact $K \subset \mathbb{R}_+^2$, we may find a rectangle $[t_1, t_2] \times [a, b]$ containing K such that $Q(t, a)$ and $Q(t, b)$ are continuous functions of t . Hence, the set

$$\{(t, y) : y(t, a) \leq y \leq y(t, b), t_1 \leq t \leq t_2\}$$

is compact and contains $T^{-1}(K)$, which is then also compact by the continuity of T .

Let $\rho = Q_{\#}\mathcal{L}^2$. Let $\tilde{\rho}$ be the density of \mathcal{L}^2 with respect to τ in \mathbb{R}_+^2 and let $\tilde{\tau}$ be the density of $\tau_{a.c.}$ with respect to \mathcal{L}^2 so that $\tilde{\rho}\tilde{\tau} = 1$, \mathcal{L}^2 -a.e. in \mathbb{R}_+^2 . The following result was proved by Wagner ([25], p.132).

Step 7. $\rho = \partial_y x = \tilde{\rho} \circ T$.

Indeed, $\rho = Q\#\mathcal{L}^2$ admits a slicing of the form $\rho = dt \otimes \tilde{\mu}_t$, where $\tilde{\mu}_t$ is the Stieltjes measure associated with the inverse of $y(t, \cdot)$, which is $x(t, y)$, and hence $\tilde{\mu}_t = \partial_y x$. Furthermore, the Lipschitz continuity of $x(t, y)$ yields

$$\rho = \partial_y x,$$

where as usual we have identified absolutely continuous measures with their densities, with respect to the corresponding Lebesgue measure. Also, we have

$$T\#\rho = T\#Q\#\mathcal{L}^2 = \mathcal{L}^2 = \tilde{\rho}\tilde{\tau} = \tilde{\rho}\tau = \tilde{\rho}T\#\mathcal{L}^2.$$

Now, for all $\phi \in C_0((0, \infty) \times \mathbb{R})$,

$$\langle \rho, \phi \rangle = \langle T\#\rho, \phi \circ Q \rangle,$$

while

$$\langle \tilde{\rho} \circ T, \phi \rangle = \langle \tilde{\rho}T\#\mathcal{L}^2, \phi \circ Q \rangle,$$

which implies

$$\rho = \tilde{\rho} \circ T,$$

where we have used that $\rho = 0$ a.e. over $T^{-1}(V)$ and V is the union of the vacuum lines, because of the first part of the assertion.

Now, let $\rho_0 = Q_0\#\mathcal{L}^1$ and $\underline{v}(t, y) = v(T(t, y))$.

Step 8. For all $\phi \in C^1_0(\mathbb{R}^2)$, we have

$$\iint_{t>0} (\rho\phi_t + \rho\underline{v}\phi_y) dt dy + \int_{t=0} \phi(0, y)\rho_0(y) dy = 0. \tag{5.5}$$

Observe that ρ and ρ_0 are uniformly bounded since $\rho = \tilde{\rho} \circ T$, $\tilde{\rho}\tilde{\tau} = 1$, and $\tilde{\tau} \geq c$ for some $c > 0$. The same holds for ρ_0 . Now, for any $\phi \in C^1_0(\mathbb{R}^2)$, we have

$$\begin{aligned} & \iint_{t>0} (\phi_t + \phi_y\underline{v})\rho dt dy + \int_{t=0} \phi(0, y)\rho_0(y) dy \\ &= \iint_{t>0} (\phi_t \circ Q + \phi_y \circ Q\underline{v} \circ Q) dt dx + \int_{t=0} \phi(0, Q_0(x)) dx \\ &= \iint_{t>0} (\phi_t \circ Q + v\phi_y \circ Q) dt dx + \int_{t=0} \phi(0, Q_0(x)) dx \\ &= \iint_{t>0} \partial_t(\phi \circ Q) dt dx + \int_{t=0} \phi(0, Q_0(x)) dx = 0, \end{aligned}$$

where we have used that $\underline{v} \circ Q = v$. In particular, we have

$$\rho_t + (\rho\underline{v})_y = 0, \tag{5.6}$$

in the sense of distributions. Now, let $x_0 \in \mathbb{R}$ be such that $y(t, x_0)$ is a Lipschitz continuous function on $[0, \infty)$ and, for a.e. t , $y(t, x_0)$ is a Lebesgue point for both $\rho(t, y)$ and $\underline{v}(t, y)$.

Step 9. There exists a solution $\bar{x}(t, y)$ of

$$\partial_y x = \rho, \quad \partial_t x = -\rho \underline{v}, \tag{5.7}$$

such that $\bar{x}(t, y(t, x_0)) = x_0$. In particular, $\bar{x}(t, y) = x(t, y)$.

Indeed, let $\rho^\varepsilon = \rho * \omega^\varepsilon$ and $(\rho \underline{v})^\varepsilon = (\rho \underline{v}) * \omega^\varepsilon$, where ω^ε is a standard mollifier. Set

$$x^\varepsilon(t, y) = \int_{y(t, x_0)}^y \rho^\varepsilon(t, s) ds + x_0.$$

Then $x^\varepsilon(t, y)$ is a Lipschitz function satisfying

$$\begin{aligned} \partial_y x^\varepsilon &= \rho^\varepsilon(t, y), \\ \partial_t x^\varepsilon &= -(\rho \underline{v})^\varepsilon(t, y) - \rho^\varepsilon(t, y(t, x_0)) \underline{v}(t, y(t, x_0)) + (\rho \underline{v})^\varepsilon(t, y(t, x_0)). \end{aligned}$$

Also,

$$|x^\varepsilon(t, y) - x_0| \leq M|y - y(t, x_0)|.$$

Hence, $x^\varepsilon(t, y)$ converges in $L^1_{loc}(\mathbb{R}^2_+)$ to a Lipschitz function $\bar{x}(t, y)$ which satisfies

$$|\bar{x}(t, y) - x_0| \leq M|y - y(t, x_0)|.$$

Thus, we have $\bar{x}(t, y(t, x_0)) = x_0$. Since

$$|\rho^\varepsilon(t, y(t, x_0)) \underline{v}(t, y(t, x_0)) - (\rho \underline{v})^\varepsilon(t, y(t, x_0))| \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0,$$

for a.e. $t > 0$, we conclude that $\bar{x}(t, y)$ satisfies (5.7). Now, since $\partial_y x(t, y) = \rho(t, y)$ and $x(t, y(t, x_0)) = x_0$, we must have $x(t, y) = \bar{x}(t, y)$.

Step 10. Equation (4.1) holds for any τ -test function $\phi(t, x)$.

Indeed, for any τ -test function $\phi(t, x)$, we have

$$\begin{aligned} & \iint_{t>0} (\phi_t \tau - v \phi_x) dt dx + \int_{t=0} \phi(0, x) \tau_0 \\ &= \iint_{t>0} (\phi_t - v \tilde{\rho} \phi_x) \tau + \int_{t=0} \phi(0, x) \tau_0 \\ &= \iint_{t>0} (\phi_t \circ T - \phi_x \circ T \rho \underline{v}) dt dy + \int_{t=0} \phi \circ T_0 dy \\ &= \iint_{t>0} \frac{\partial}{\partial t} (\phi \circ T) dt dy + \int_{t=0} \phi \circ T_0 dy \\ &= - \int_{\mathbb{R}} \lim_{\delta \rightarrow 0} \phi \circ T(y, \delta) dy + \int_{t=0} \phi \circ T_0 dy \\ &= 0. \end{aligned}$$

□

Acknowledgements. Gui-Qiang Chen’s research was supported in part by the National Science Foundation under Grants DMS-0204225, DMS-0204455, and INT-9987378. Hermano Frid’s research was supported in part by CNPq through the grants 352871/96-2, 465714/00-5, 479416/01-0 and FAPERJ through the grant E-26/151.890/2000.

References

1. Anzellotti, G.: Pairings between measures and functions and compensated compactness. *Ann. Mat. Pura Appl.* **135**, 293–318 (1983)
2. Baiocchi, C., Capelo, A.: *Variational and Quasi-Variational Inequalities, Applications to Free-Boundary Problems*. Vols. **1, 2**, Chichester-New York: Wiley, 1984
3. Bianchini, S., Bressan, A.: *Vanishing viscosity solutions of nonlinear hyperbolic systems*. Preprint, SISSA, Italy, 2001
4. Brezzi, F., Fortin, M.: *Mixed and Hybrid Finite Element Methods*. New York: Springer-Verlag, 1991
5. Chen, G.-Q., Frid, H.: Divergence-measure fields and conservation laws. *Arch. Rat. Mech. Anal.* **147**, 89–118 (1999)
6. Chen, G.-Q., Frid, H.: Decay of entropy solutions of nonlinear conservation laws. *Arch. Rat. Mech. Anal.* **146**, 95–127 (1999)
7. Chen, G.-Q., Frid, H.: Large-time behavior of entropy solutions of conservation laws. *J. Diff. Eqs.* **152**, 308–357 (1999)
8. Chen, G.-Q., Frid, H.: Uniqueness and asymptotic stability of Riemann solutions for the compressible Euler equations. *Trans. Am. Math. Soc.* **353**, 1103–1117 (2001)
9. Chen, G.-Q., Frid, H., Li, Y.: Uniqueness and stability of Riemann solutions with large oscillation in gas dynamics. *Commun. Math. Phys.* **228**, 201–217 (2002)
10. Dafermos, C.M.: *Hyperbolic Conservation Laws in Continuum Physics*. Berlin: Springer-Verlag, 1999
11. DiPerna, R.: Uniqueness of solutions to hyperbolic conservation laws. *Indiana Univ. Math. J.* **28**, 137–188 (1979)
12. Evans, L.C., Gariepy, R.F.: *Lecture Notes on Measure Theory and Fine Properties of Functions*. Boca Raton, Florida: CRC Press, 1992
13. Federer, H.: *Geometric Measure Theory*. Berlin-Heidelberg-New York: Springer-Verlag, 1969
14. Gagliardo, E.: Caratterizzazioni delle tracce sulla frontiera relativa ad alcune classi di funzioni in n variabili. *Rend. Sem. Mat. Univ. Padova* **27**, 284–305 (1957)
15. Glimm, J.: Solutions in the large for nonlinear hyperbolic systems of equations. *Commun. Pure Appl. Math.* **18**, 95–105 (1965)
16. Lax, P.D.: *Hyperbolic Systems of Conservation Laws and the Mathematical Theory of Shock Waves*. CBMS. **11**, Philadelphia: SIAM, 1973
17. Lax, P.D.: Shock waves and entropy. In: *Contributions to Functional Analysis*. E.A. Zarantonello (ed). New York: Academic Press, 1971, pp. 603–634
18. Liu, T.-P., Smoller, J.: On the vacuum state for the isentropic gas dynamics equations. *Adv. Appl. Math.* **1**, 345–359 (1980)
19. Perthame, B.: *Kinetic Formulation of Conservation Laws*. Lecture Notes, Ecole Normale Supérieure, Paris, 2001
20. Rodrigues, J.-F.: *Obstacle Problems in Mathematical Physics*. North-Holland Mathematics Studies **134**, Amsterdam: Elsevier Science Publishers B.V., 1987
21. Schwartz, L.: *Théorie des Distributions* (2 volumes). Actualités Scientifiques et Industrielles **1091, 1122**, Paris: Herman, 1950–51
22. Serre, D.: *Systems of Conservation Laws*. Vols. **1, 2**, Cambridge: Cambridge University Press, 1999, 2000
23. Stein, E.: *Singular Integrals and Differentiability Properties of Functions*. Princeton, NJ: Princeton Univ. Press, 1970
24. Volpert, A.I.: The space BV and quasilinear equations. *Mat. Sb. (N.S.)* **73**, 255–302 (1967) *Math. USSR Sbornik* **2**, 225–267 (1967) (in English)
25. Wagner, D.: Equivalence of the Euler and Lagrange equations of gas dynamics for weak solutions. *J. Diff. Eqs.* **68**, 118–136 (1987)
26. Whitney, H.: *Geometric Integration Theory*. Princeton, NJ: Princeton Univ. Press, 1957
27. Whitney, H.: Analytic extensions of differentiable functions defined in closed sets. *Trans. Am. Math. Soc.* **36**, 63–89 (1934)
28. Ziemer, W.P.: Cauchy flux and sets of finite perimeter. *Arch. Rational Mech. Anal.* **84**, 189–201 (1983)
29. Ziemer, W.P.: *Weakly Differentiable Functions: Sobolev Spaces and Functions of Bounded Variation*. New York: Springer-Verlag, 1989