

POINCARÉ RECURRENCE AND NUMBER THEORY: THIRTY YEARS LATER

BRYNA KRA

Hillel Furstenberg's 1981 article in the *Bulletin* gives an elegant introduction to the interplay between dynamics and number theory, summarizing the major developments that occurred in the few years after his landmark paper [21]. The field has evolved over the past thirty years, with major advances on the structural analysis of dynamical systems and new results in combinatorics and number theory. Furstenberg's article continues to be a beautiful introduction to the subject, drawing together ideas from seemingly distant fields.

Furstenberg's article [21] gave a general correspondence between regularity properties of subsets of the integers and recurrence properties in dynamical systems, now dubbed the *Furstenberg Correspondence Principle*. He then showed that such recurrence properties always hold, proving what is now referred to as the *Multiple Recurrence Theorem*. Combined, these results gave a new proof of Szemerédi's Theorem [45]: if $S \subset \mathbb{Z}$ has positive upper density, then S contains arbitrarily long arithmetic progressions. This proof led to an explosion of activity in ergodic theory and topological dynamics, beginning with new proofs of classic results of Ramsey Theory and ultimately leading to significant new combinatorial and number theoretic results. The full implications of these connections have yet to be understood.

The approach harks back to the earliest results on recurrence, in the measurable setting and in the topological setting. A measure preserving system is a quadruple (X, \mathcal{B}, μ, T) , where X denotes a set, \mathcal{B} is a σ -algebra on X , μ is a probability measure on (X, \mathcal{B}) , and $T: X \rightarrow X$ is a measurable transformation such that $\mu(T^{-1}(A)) = \mu(A)$ for all $A \in \mathcal{B}$. Poincaré Recurrence states that if (X, \mathcal{B}, μ, T) is a measure preserving system and $A \in \mathcal{B}$ with $\mu(A) > 0$, there exists $n \in \mathbb{N}$ such that $\mu(A \cap T^{-n}A) > 0$. A (topological) dynamical system is a pair (X, T) , where X is a compact metric space and $T: X \rightarrow X$ is a continuous map. One can show that any such topological space admits a Borel, probability measure that preserves T . In particular, Poincaré Recurrence implies recurrence in the topological setting. Birkhoff [13] gave a direct proof of

The author was partially supported by NSF grant 0900873.

this, showing that in any dynamical system (X, T) , there exists $x \in X$ such that $T^{n_k}x \rightarrow x$ for some sequence of integers $n_k \rightarrow \infty$.

Birkhoff's recurrence was generalized by Furstenberg and Weiss [26], who showed that given a dynamical system (X, T) and an integer $\ell \geq 1$, there exists $x \in X$ such that $T^{n_k}x \rightarrow x, T^{2n_k}x \rightarrow x, \dots, T^{\ell n_k}x \rightarrow x$ for some sequence of integers $n_k \rightarrow \infty$. Using an analog of the Furstenberg Correspondence Principle adapted to topological dynamical systems, this in turn implies van der Waerden's Theorem: in any finite partition of the integers, some piece contains arbitrarily long arithmetic progressions. Topological methods were then used to prove numerous other partition results, including Schur's Theorem, a multidimensional version of van der Waerden's Theorem, and the Hales-Jewett Theorem. As a sample of these techniques, Furstenberg gives a proof of a polynomial result, shown independently by Sárközy [44]: in any finite partition of the integers, some piece contains two integers which differ by a square. Vast generalizations of such polynomial results were given by Bergelson and Leibman ([7] and [8]).

In the measure theoretic setting, Furstenberg proved a far reaching generalization of the Poincaré Recurrence Theorem in his Multiple Recurrence Theorem: if (X, \mathcal{B}, μ, T) is a measure preserving system, $\ell \geq 1$ is an integer, and $A \in \mathcal{B}$ has positive measure, then there exists $n \in \mathbb{N}$ such that

$$(1) \quad \mu(A \cap T^{-n}A \cap T^{-2n}A \cap \dots \cap T^{-\ell n}A) > 0.$$

The study of which iterates n are possible has received significant attention, with polynomial iterates [7], generalized polynomials [12], sequences arising from Hardy-Fields [17], shifts of the primes ([?] and [18]), and this result has been generalized in numerous other ways (see for example [25] and [40]). Many of these results have yet to be proved using methods that do not rely on dynamics.

Via the Furstenberg Correspondence Principle, the analog of Equation (1) with the transformations T, T^2, \dots, T^ℓ replaced by commuting transformations T_1, T_2, \dots, T_ℓ leads to a multidimensional Szemerédi Theorem, and this and generalizations were proven by Furstenberg and Katznelson ([23] and [24]). This too has been studied further, including restrictions on iterates (see [7], [11], and [19]) and generalizations to other groups (for example [41]). Again, many of these results have yet to be proven via combinatorial methods.

Using ergodic theory, the natural approach to prove positivity of an expression such as that in Equation (1) is to take the average for $1 \leq n \leq N$ and show that the liminf of this average is positive as

$N \rightarrow \infty$. More generally, one can consider convergence of averages

$$(2) \quad \frac{1}{N} \sum_{n=1}^N f_1(T_1^{a_1(n)}x) f_2(T_2^{a_2(n)}x) \dots f_\ell(T_\ell^{a_\ell(n)}x),$$

where (X, \mathcal{B}, μ) is a probability space, $T_1, T_2, \dots, T_\ell: X \rightarrow X$ are commuting, measure preserving transformations, $f_1, f_2, \dots, f_\ell \in L^\infty(\mu)$, and the exponents $a_1(n), a_2(n), \dots, a_\ell(n)$ are sequences of integers. The existence of the limit in $L^2(\mu)$, and the study of the structures controlling the limiting behavior of such averages, has received much attention within ergodic theory and has more recently lead to new number theoretic results. The case when all the transformations T_i are equal with linear exponents is fully understood, with partial results, for example in [21] and [15], and the complete convergence in [32]. These results have been generalized and viewed in other ways, with further studies of the linear case [49], polynomial iterates ([27], [33] and [41]), commuting transformations ([16] and [46]), restrictions on the iterates for commuting transformations (see for example [36], [14], [1], and [2]), nilpotent group actions [9], and the corresponding average for flows (see [43], [10] and [3]). For a single transformation, we have a complete understanding in [32] of the structures controlling convergence (with a topological analog in [35]). The general convergence and associated structures controlling such averages are yet to be understood.

The connections to number theoretic and combinatorial problems continue to grow, particularly with the spectacular breakthrough of Green and Tao [28] showing that the primes contain arbitrarily long arithmetic progressions. While there is no explicit use of ergodic theory in Green and Tao's proof, the methods used by Furstenberg [21] influence the approach. In more recent work on asymptotics of the number of progressions in the primes ([29], [30], and [31]) and other connections between ergodic theory and number theory (see, for example [6], [34], and [20]), the structures controlling the averages of the form in Equation (2) play a prominent role.

This is only a brief overview of current areas of research with origins in the work surveyed in Furstenberg's *Bulletin* article. An extensive introduction to the field is contained in Furstenberg's book [22]. There are more recent surveys on recurrence, ergodic Ramsey Theory, convergence problems, and connections to number theory (see for example [4], [5], [42], [37], [38], [39], [47], and [48]), that are natural continuations of the topics reviewed in the accompanying article.

REFERENCES

- [1] T. Austin. Pleasant extensions retaining algebraic structure, I. arXiv:0905.0518
- [2] T. Austin. Pleasant extensions retaining algebraic structure, II. arXiv:0910.0907
- [3] T. Austin. Norm convergence of continuous-time polynomial multiple ergodic averages. arXiv:1103.0223v2
- [4] V. Bergelson. Combinatorial and Diophantine Applications of Ergodic Theory (with appendices by A. Leibman and by A. Quas and M. Wierdl). *Handbook of Dynamical Systems*, Vol. 1B, (Eds.: B. Hasselblatt, A. Katok). Elsevier (2006), 745–841.
- [5] V. Bergelson. Ergodic Ramsey theory an update. *Ergodic Theory of \mathbb{Z}^d -actions*, (Eds.: M. Pollicott, K. Schmidt). Cambridge University Press, Cambridge (1996), 1–61.
- [6] V. Bergelson, B. Host, B. Kra, with an appendix by I. Ruzsa. Multiple recurrence and nilsequences. *Invent. Math.*, **160** (2005), 261–303.
- [7] V. Bergelson, A. Leibman. Polynomial extensions of van der Waerden’s and Szemerédi’s theorems. *Journal of AMS* **9** (1996), 725–753.
- [8] V. Bergelson, A. Leibman. Set-polynomials and polynomial extension of the Hales-Jewett theorem. *Ann. of Math.* **150** (1999), 33–75.
- [9] V. Bergelson, A. Leibman. A nilpotent Roth theorem. *Invent. Math.* **147** (2002), 429–470.
- [10] V. Bergelson, A. Leibman, C. G. Moreira. From discrete to continuous time ergodic theorems. Preprint.
- [11] V. Bergelson, A. Leibman, T. Ziegler. The shifted primes and the multidimensional Szemerédi and polynomial van der Waerden theorems. To appear, *Comptes Rendus Math.*
- [12] V. Bergelson, R. McCutcheon. Idempotent ultrafilters, multiple weak mixing and Szemerédi’s theorem for generalized polynomials. *J. Analyse Math.* **111** (2010), 77–130.
- [13] G. D. Birkhoff. *Dynamical Systems*. Amer. Math. Soc. Colloq. Publ. **9** American Mathematical Society, Providence, 1927.
- [14] Q. Chu, N. Frantzikinakis, B. Host. Ergodic averages of commuting transformations with distinct degree polynomial iterates. To appear, *Proc. Lond. Math. Soc.*
- [15] J. -P. Conze, E. Lesigne. Théorèmes ergodiques pour des mesures diagonales. *Bull. Soc. Math. France* **112** (1984), 143–175.
- [16] J. -P. Conze, E. Lesigne. Sur un théorème ergodique pour des mesures diagonales. *Publ. Inst. Rech. Math. Rennes 1987-1* (1988), 1–31.
- [17] N. Frantzikinakis. Multiple recurrence and convergence for Hardy sequences of polynomial growth. *J. Analyse Math.* **112** (2010), 79–135.
- [18] N. Frantzikinakis, B. Host, B. Kra. Multiple recurrence and convergence for sets related to the primes. *J. Reine Angew. Math.*, **611** (2007), 131–144.
- [19] N. Frantzikinakis, B. Host, B. Kra. The polynomial multidimensional Szemerédi Theorem along shifted primes. arXiv:1009.1484
- [20] N. Frantzikinakis, E. Lesigne, M. Wierdl. Powers of sequences and recurrence. *Proc. London. Math. Soc.*, **98** (2009), 504–530.

- [21] H. Furstenberg. Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions. *Journal d'Analyse Math.* **31** (1977), 204–256.
- [22] H. Furstenberg. Recurrence in Ergodic Theory and Combinatorial Number Theory. Princeton University Press, Princeton, 1981.
- [23] H. Furstenberg, Y. Katznelson. An ergodic Szemerédi theorem for commuting transformations. *Journal d'Analyse Math.* **34** (1978), 275–291.
- [24] H. Furstenberg, Y. Katznelson. An ergodic Szemerédi theorem for IP-systems and combinatorial theory. *Journal d'Analyse Math.* **45** (1985), 117–168
- [25] H. Furstenberg, Y. Katznelson. A density version of the Hales-Jewett theorem. *Journal d'Analyse Math.* **57** (1991), 64–119.
- [26] H. Furstenberg, B. Weiss. Topological dynamics and combinatorial number theory. *Journal d'Analyse Math.* **34** (1978), 61–85.
- [27] H. Furstenberg, B. Weiss. A mean ergodic theorem for $\frac{1}{N} \sum_{n=1}^N f(T^n x)g(T^{n^2} x)$. Convergence in Ergodic Theory and Probability (Columbus, OH, 1993), Walter de Gruyter, 1996, 193–227.
- [28] B. Green, T. Tao. The primes contain arbitrarily long arithmetic progressions. *Ann. of Math.*, **167** (2008), 481–547.
- [29] B. Green, T. Tao. Linear Equations in the Primes. *Ann. of Math.*, **171** (2010), 1753–1850.
- [30] B. Green, T. Tao. The Möbius function is strongly orthogonal to nilsequences. To appear, *Ann. of Math.*
- [31] B. Green, T. Tao, T. Ziegler. An inverse theorem for the Gowers U^k norm. arxiv:1009.3998.
- [32] B. Host, B. Kra. Nonconventional averages and nilmanifolds. *Ann. of Math.*, **161** (2005), 398–488.
- [33] B. Host, B. Kra. Convergence of Polynomial ergodic averages. *Isr. J. Math.*, **149** (2005), 1–19.
- [34] B. Host, B. Kra. Uniformity norms on ℓ^∞ and applications. *J. d'Analyse Mathématique*, **108** (2009), 219–276.
- [35] B. Host, B. Kra, A. Maass. Nilsequences and a structure theorem for topological dynamical systems. *Adv. in Math.*, **224** (2010), 103–129.
- [36] M. Johnson. Convergence of Polynomial Ergodic Averages. *Illinois J. Math.*, **53** (2009), 865–882.
- [37] B. Kra. The Green-Tao Theorem on arithmetic progressions in the primes: an ergodic point of view. *Bull. Amer. Math. Soc.*, **43** (2006), 3–23.
- [38] B. Kra. From combinatorics to ergodic theory and back again. *Proceedings of the International Congress of Mathematicians*, Madrid 2006, Vol. III, 57–76.
- [39] B. Kra. Ergodic methods in additive combinatorics. Additive combinatorics, *CRM Proc. Lecture Notes* **43**, Amer. Math. Soc., Providence, RI (2007), 103–144.
- [40] A. Leibman. Multiple recurrence theorem for measure preserving actions of a nilpotent group. *Geom. and Funct. Anal.*, **8** (1998), 853–931.
- [41] A. Leibman. Convergence of multiple ergodic averages along polynomials of several variables. *Isr. J. Math.*, **146** (2005), 303–316.

- [42] R. McCutcheon. Elementary methods in ergodic Ramsey theory. Lecture Notes in Mathematics 1722. Springer-Verlag, Berlin, 1999.
- [43] A. Potts. Multiple ergodic averages for flows and an application. To appear, *Illinois J. Math.*
- [44] On difference sets of integers III. *Acta Math. Acad. Sci. Hungar.*, **31** (1978), 125–149.
- [45] E. Szemerédi. On sets of integers containing no k elements in arithmetic progression. *Acta Arith.* **27** (1975) 199–245.
- [46] T. Tao. Norm convergence of multiple ergodic averages for commuting transformations. *Erg. Th. Dynam. Sys.*, **28** (2008), 657–688.
- [47] T. Tao. The ergodic and combinatorial approaches to Szemerédi’s theorem. Additive combinatorics, *CRM Proc. Lecture Notes* **43**, Amer. Math. Soc., Providence, RI (2007), 145–193.
- [48] T. Tao. The dichotomy between structure and randomness, arithmetic progressions, and the primes. *Proceedings of the International Congress of Mathematicians*, Madrid 2006, Vol. I, 581–608.
- [49] T. Ziegler. Universal characteristic factors and Furstenberg averages. *J. Amer. Math. Soc.*, **20** (2007), 53–97.

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, 2033 SHERIDAN ROAD, EVANSTON, IL 60208-2730, USA
E-mail address: `kra@math.northwestern.edu`